


A Practical Handbook for Determining the Ages of Gulf of Mexico Fishes

Second Edition



Gulf States Marine Fisheries Commission

Publication Number 167

July 2009

A Practical Handbook for Determining the Ages of Gulf of Mexico Fishes

Second Edition

edited by:

Steve VanderKooy

**Gulf States Marine Fisheries Commission
2404 Government St
Ocean Springs, MS 39564**

Publication Number 167

July 2009

A publication of the Gulf States Marine Fisheries Commission pursuant to National Oceanic and Atmospheric Administration Award Number NA05NMF4070005. This paper is funded by a grant from the National Oceanic and Atmospheric Administration. The views expressed herein are those of the author(s) and do not necessarily reflect the views of NOAA or any of its subagencies.



Contributors

Ms. Janet Tunnell	Fish and Wildlife Research Institute/FWC
Dr. Katherine Guindon	Fish and Wildlife Research Institute/FWC
Mr. Ken Edds	Louisiana Department of Wildlife and Fisheries
Ms. Isis Longo	Louisiana Department of Wildlife and Fisheries
Dr. Gary Fitzhugh	NOAA Fisheries/Panama City Laboratory
Mr. Andrew J. Fischer	Louisiana State University
Dr. Luiz Barbieri	Fish and Wildlife Research Institute/FWC
Dr. Robert Allman	NOAA Fisheries /Panama City Laboratory
Mr. Doug DeVries	NOAA Fisheries /Panama City Laboratory
Mr. Chris Palmer	NOAA Fisheries /Panama City Laboratory
Dr. Mike Murphy	Fish and Wildlife Research Institute/Florida FWC
Dr. Debra Murie	University of Florida
Ms. Carrie Lee Fioramonti	NOAA Fisheries /Panama City Laboratory
Ms. Julia Clifton	Louisiana Department of Wildlife and Fisheries
Mr. Read Hendon	USM/CMS/Gulf Coast Research Laboratory
Mr. Erick Porche	Mississippi Department of Marine Resources
Mr. Britt Bumguardner	Texas Parks and Wildlife Department
Mr. Bob Colura	Texas Parks and Wildlife Department
Mr. Dan Merryman	Fish and Wildlife Research Institute/FWC
Mr. Eric Robillard	Fish and Wildlife Research Institute/FWC
Mr. John Mareska	ADCNR /Marine Resources Division
Dr. Richard McBride	Fish and Wildlife Research Institute/FWC
Mr. Wesley Devers	Mississippi Department of Marine Resources
Mr. Joseph W. Smith	NOAA Fisheries /Beaufort Laboratory
Dr. Walter Ingram	NOAA Fisheries /Pascagoula Laboratory
Ms. Beverly Barnett	NOAA Fisheries /Panama City Laboratory
Mr. Gary Gray	USM/CMS/Gulf Coast Research Laboratory
Dr. Stephen Wischniowski	Old Dominion University
Dr. Stephen Bobko	Old Dominion University
Ms. Jaime Miller	ADCNR /Marine Resources Division
Ms. Brandy Coates	ADCNR /Marine Resources Division
Mr. John R. Foster	Georgia Department of Natural Resources
Ms. Gabe Gaddis	Georgia Department of Natural Resources
Mr. Jim Duffy	ADCNR/Marine Resources Division
Ms. Linda Lombardi-Carlson	NOAA Fisheries/Panama City Laboratory
Ms. Ivy Baremore	NOAA Fisheries/Panama City Laboratory
Ms. Donna McDowell	Georgia Department of Natural Resources
Mr. James “Tut” Warren	USM/CMS/Gulf Coast Research Laboratory

GSMFC Staff

Mr. Larry B. Simpson
Executive Director

Mr. Steven J. VanderKooy
Program Coordinator

Mrs. Teri L. Freitas
Staff Assistant

Gulf States Marine Fisheries Commission

Commissioners and Proxies

ALABAMA

Barnett Lawley, Commissioner
Alabama Department of Conservation
and Natural Resources
64 North Union Street
Montgomery, AL 36130-1901
Proxy:
Vernon Minton, Director
Alabama Marine Resources Division
P.O. Drawer 458
Gulf Shores, AL 36547

Representative Spencer Collier
P.O. Box 550
Irvington, AL 36544

Chris Nelson
Bon Secour Fisheries, Inc.
P.O. Box 60
Bon Secour, AL 36511

FLORIDA

Ken Haddad, Executive Director
Florida Fish and Wildlife Conservation Commission
620 South Meridian Street
Tallahassee, FL 32399-1600
Proxy:
Virginia Vail
FWC Division of Marine Fisheries
620 South Meridian Street
Tallahassee, FL 32399-1600

Senator Thad Altman
State Senator, District 24
6767 North Wickham Road, Suite 211
Melbourne, FL 32940

Hayden R. Dempsey
Greenberg Traurig, P.A.
P.O. Box 1838
Tallahassee, FL 32302

LOUISIANA

Robert Barham, Secretary
Louisiana Department of Wildlife and Fisheries
P.O. Box 98000
Baton Rouge, LA 70898-9000
Proxy:
Randy Pausina
Louisiana Department of Wildlife and Fisheries
P.O. Box 98000
Baton Rouge, LA 70898-9000

Senator Butch Gautreaux, *Chairman*
714 2nd Street
Morgan City, LA 70380

Wilson Gaidry
8911 Park Avenue
Houma, LA 70363

MISSISSIPPI

William Walker, Executive Director
Mississippi Department of Marine Resources
1141 Bayview Avenue, Suite 101
Biloxi, MS 39530
Proxy:
Dale Diaz
Mississippi Department of Marine Resources
1141 Bayview Avenue
Biloxi, MS 39530

Senator Tommy Gollott
235 Bay View Avenue
Biloxi, MS 39530

Joe Gill Jr.
Joe Gill Consulting, LLC
P.O. Box 535
Ocean Springs, MS 39566-0535

TEXAS

Carter Smith, Executive Director
Texas Parks and Wildlife Department
4200 Smith School Road
Austin, TX 78744
Proxy:
Mike Ray
Coastal Fisheries Division
Texas Parks and Wildlife Department
4200 Smith School Road
Austin, TX 78744

Senator Mike Jackson
Texas Senate
P.O. Box 12068
Austin, TX 78711

David McKinney
44 East Avenue, Suite 315
Austin, TX 78701

Table of Contents

Contributors	ii
Commissioners and Proxies	iii
Table of Contents	iv
Abbreviations	vi
Acknowledgments	viii
Preface	viii
1.0 Introduction	1-1
2.0 Otolith Structure and Function	2-1
3.0 General Processing Techniques	3-1
3.1 Otolith Removal	3-1
3.2 Cleaning and Storage	3-3
3.3 Sectioning Preparation	3-3
3.3.1 Embedding Otoliths	3-4
3.3.1.1 Embedding Whole Otoliths	3-4
3.3.1.2 Embedding Small Otoliths in Bullet Molds	3-5
3.3.1.3 Marking the Core	3-5
3.3.2 Mounting Whole Otoliths on Slides	3-6
3.3.2.1 Marking the Core	3-6
3.3.3 Thin Sectioning Machine Preparation	3-7
3.4 Sectioning Techniques	3-7
3.4.1 High Speed Wafering Saw	3-7
3.4.1.1 Embedded Otoliths	3-8
3.4.1.2 Whole Otoliths	3-8
3.4.2 Low Speed Wafering Saw	3-9
3.4.2.1 Embedded Otoliths	3-9
3.4.2.2 Whole Mounted Otoliths	3-10
3.4.2.2.1 Production Ageing	3-12
3.4.3 Thin Section Machine	3-12
3.5 Mounting Sections	3-14
3.6 Alternative Techniques	3-15
3.6.1 Whole Otoliths	3-15
3.6.2 Break and Burn	3-16
3.6.2.1 Reading and Interpretation	3-16
3.6.3 Scales	3-17
3.6.3.1 Scale Collection and Handling	3-17
3.6.3.1.1 Raw Scales	3-17
3.6.3.1.2 Scale Impressions	3-18
3.6.3.2 Scale Reading and Interpretation	3-19
3.6.4 Spines and Finrays	3-21
3.6.4.1 Sectioning Spines	3-22
3.6.4.1.1 Thin Section Machine	3-22
3.6.4.2.2 Low Speed Wafering Saw	3-23
3.6.4.2 Sectioning Finrays	3-23
3.6.4.3 Reading and Interpretation	3-24
3.6.5 Vertebrae	3-25
3.6.5.1 Extraction and Storage	3-25
3.6.5.2 Sectioning and Reading	3-26

3.7	Section Enhancement.....	3-26
3.7.1	Polishing	3-26
3.7.2	Etching	3-27
3.7.3	Staining	3-28
3.7.4	Clearing.....	3-28
3.7.5	Baking.....	3-28
3.7.6	Filters	3-29
3.8	Microscopy, Image Analysis, and Measurements.....	3-29
4.0	Age Determination	4-1
4.1	Otolith Development.....	4-1
4.2	Ring Enumeration	4-2
4.2.1	Margin Codes.....	4-3
4.2.1.1	Quality of Otolith Section Margins.....	4-4
4.3	Assignment of Age	4-4
4.3.1	Biological Age	4-6
4.3.2	Assigning Year Groups.....	4-7
4.4	Quality Control in Processing	4-7
4.4.1	Validation.....	4-8
4.4.1.1	Chemical Marking	4-8
4.4.1.2	Marginal Increment Analysis.....	4-8
4.4.2	Accuracy	4-9
4.4.3	Precision.....	4-9
4.4.4	Reference Collection.....	4-9
4.4.5	Reader Comparisons	4-10
4.5	Other Parameters and Their Usefulness.....	4-11
5.0	Species-Specific Otolith Characteristics and Processing Details.....	5-1
5.1	Red Drum <i>Sciaenops ocellatus</i>	5-2
5.2	Spotted Seatrout <i>Cynoscion nebulosus</i>	5-7
5.3	Black Drum <i>Pogonias cromis</i>	5-11
5.4	Striped Mullet <i>Mugil cephalus</i>	5-15
5.5	Southern Flounder <i>Paralichthys lethostigma</i>	5-19
5.6	Gray Triggerfish <i>Balistes caprisacus</i>	5-23
5.7	Red Snapper <i>Lutjanus campechanus</i>	5-27
5.8	King Mackerel <i>Scomberomorus cavalla</i>	5-31
5.9	Greater Amberjack <i>Seriola dumerili</i>	5-36
5.10	Spanish Mackerel <i>Scomberomorus maculatus</i>	5-41
5.11	Atlantic Croaker <i>Micropogonias undulatus</i>	5-45
5.12	Sheepshead <i>Archosargus probatocephalus</i>	5-50
5.13	Gag <i>Mycteroperca microlepis</i>	5-54
5.14	Red Grouper <i>Epinephelus morio</i>	5-58
5.15	Gray Snapper <i>Lutjanus griseus</i>	5-62
5.16	Vermilion Snapper <i>Rhomboplites aurorubens</i>	5-66
5.17	Gulf Menhaden <i>Brevoortia patronus</i>	5-70
6.0	Literature Cited	6-1
7.0	Glossary	7-1
8.0	Appendices.....	8-1
8.1	Suppliers and Supplies - Permanent Equipment and Apparatus.....	8-2
8.2	Suppliers and Supplies - Consumables	8-4
8.3	Suppliers and Supplies - Supplier Contact Information.....	8-9
8.4	Photo Credits.....	8-11

Abbreviations and Symbols

ADCNR/MRD	Alabama Department of Conservation and Natural Resources/Marine Resources Division
DMS	Data Management Subcommittee
FWC	Florida Fish and Wildlife Conservation Commission
FWRI	Fish and Wildlife Research Institute
FIN	Fishery Information Network
g	gram
GCRL	Gulf Coast Research Laboratory
GSI	gonadosomatic index
GMFMC	Gulf of Mexico Fisheries Management Council
GSMFC	Gulf States Marine Fisheries Commission
IJF	interjurisdictional fisheries
kg	kilogram
km	kilometer
lbs	pounds
LDWF	Louisiana Department of Wildlife and Fisheries
LSU	Louisiana State University
m	meter
MIA	marginal increment analysis
mm	millimeters
MDMR	Mississippi Department of Marine Resources
MFCMA	Magnuson Fishery Conservation and Management Act
MRFSS	Marine Recreational Fisheries Statistics Survey
MSDS	material safety data sheets
n	number or sample size
NMFS	National Marine Fisheries Service
SAT	Stock Assessment Team
SD	standard deviation
SE	standard error
SEM	scanning electron microscope
SL	standard length
TL	total length
TPWD	Texas Parks and Wildlife Department
TW	total weight
VIMS	Virginia Institute of Marine Sciences

Acknowledgments

The Gulf States Marine Fisheries Commission would like to thank all those who provided input, support, materials, and review for the Second Edition of the manual. Thanks go to the past and present staff at the Gulf Coast Research Laboratory in Ocean Springs Mississippi, the FWRI in St. Petersburg Florida, and the NMFS Panama City Laboratory who provided specimens and time for many of the photographic contributions. We would especially like to thank Dr. Leonard Maiolatesi and Steve Perrigin, captains of the charter boat the “Fighting Chicken” and “Strictly Fishing”, and Carlos Llull of the FWRI in Pensacola, Florida for the assistance in collecting fish heads for the new species sections. Finally, special thanks go out again to Dr. Stacey Randall, DVM, and the staff at Bienville Animal Medical Center in Ocean Springs, Mississippi, for putting up with numerous severed, frozen fish heads in our effort to generate the detailed x-rays used throughout this manual.

Preface

In 1995, the Stock Assessment Team (SAT) of the Gulf States Marine Fisheries Commission (GSMFC) proposed a manual to facilitate consistent, quality age determination of exploited Gulf of Mexico fishes and outline methodologies employed by the Gulf's marine agencies to process the hard parts. The SAT recognized that its charge to integrate state-specific stock assessments for GSMFC fishery management plans would require consistent criteria for age determinations of fishes throughout the Gulf. Therefore, a work group of experienced fisheries professionals was assembled to develop and expand this manual. The work group is comprised of two individuals from each state agency along with contributors from academia and the National Oceanographic and Atmospheric Administration's (NOAA's) National Marine Fisheries Service (NMFS). The original 'otolith manual' was completed in 2003 after two years of effort by the work group. The revision to the manual was begun almost immediately after its initial printing but due to a number of events such as hurricanes Katrina and Rita in 2005 and funding issues, the revision was delayed.

The Second Edition is a continuation of the standardization effort developed previously by the Gulf States Marine Fisheries Commission's Otolith Work Group. This revision has contributions from a number of individuals from the original publication. In total, fourteen state agencies, federal laboratories, and universities provided material and/or reviews for this edition. Again, because the majority of fish ages in the Gulf states are determined by otolith interpretation, this manual focuses primarily on otoliths although more time is dedicated to the alternate techniques such as scales and fin spines. This edition includes several additional species for which data are currently being collected under the Gulf's Fisheries Information Network, or FIN Program. Those FIN species have been identified for stock assessment in NOAA's Southeast Data, Assessment, and Review (SEDAR) process in the near future. Standardization of the techniques for them should facilitate a smoother assessment process. In addition, several new or improved processing techniques have been added or modified from the original descriptions.

We have tried to provide information on all the various techniques that have proven to be useful or unsuccessful for each of the species covered in Section 5.0 and we have updated each species ensuring that validation is included where possible. As always, when new species are added to the manual in the future, these techniques will be expanded where appropriate. These additions will be available on-line at the GSMFC website (www.gsmfc.org) or through the Gulf States Marine Fisheries Commission office.

1.0 Introduction

Fisheries science has been at the forefront of studies on animal growth and population dynamics in part because the age of individual fish can be determined. The original technique used for estimating ages of fishes involved following modal progressions of fish lengths as they changed through time (Petersen 1892). Later, marks on the animal's hard parts (calcified structures) were found to be formed on a regular and sometimes annual basis (Hoffbauer 1898, Reibisch 1899, Heinke 1905). These hard parts include scales, bones, spines, vertebrae, and otoliths. Of these, otoliths appear to be the least sensitive to changes in fish condition (Campana and Neilson 1985). Otolith growth is allometric and enough material is continuously deposited on its medial surface that marks in the form of rings are distinguishable throughout the life of most fishes. This provides a reliable and permanent record of temporal features.

The significance of determining age is that it allows fishery scientists to relate their observations to a time frame and estimate various biological rates by species. Ages of individual fish are required to estimate growth rate, age at recruitment, maturity schedules, and age-specific fecundity for a specific species. In addition, the calculation of natural and fishing mortality rates and age-specific sex ratios also require age data. In the simplest sense, this time frame may involve estimating the number of years a fish spends in a particular life history stage or habitat or determining the number of years that fishes are available for harvest.

Age determination has become such an integral part of the analyses of exploited fish populations that most agencies responsible for fisheries management have begun to routinely collect and process otoliths taken from fish sampled using fishery-dependent and fishery-independent methods. The technical skills and

equipment needed for 'production ageing' are variable depending on the type of fish and the objectives of the study.

Numerous publications have been written that describe these techniques for sampling, processing, and analyzing otoliths for age determination. Pentilla and Dery (1988) documented age determination techniques used by the staff at the Woods Hole Laboratory, National Marine Fisheries Service (NMFS) to process samples from Northwest Atlantic fishes and mollusks. Other reports have targeted the interpretation of daily growth increments (Pannella 1971) from larval and juvenile fishes using equipment and techniques similar to those used for adult fishes (Secor et al. 1991, Stevenson and Campana 1992). In addition, the use of otoliths as records of age, stock identification, pollution exposure, and various environmental conditions during the life of a fish has developed into an inter-disciplinary scientific field (Secor et al. 1995a).

Recent advances in microchemistry with regard to the use of diagnostic isotopes incorporated within the aragonite matrix of the otolith allows scientists to evaluate extant populations and the area or environment they grew in as well as the environmental conditions of long extinct populations. These advanced techniques are not detailed in this manual since its focus is on the processing of hardparts for direct, visual interpretation. However, a number of papers describing the techniques have been published (Campana et al. 1995b and 1997, Sinclair et al. 1998, Radtke et al. 1999, Markwitz et al. 2000, Limburg et al. 2007) along with a thorough review of the most common methodologies, their assumptions, and their value as life history tools (Elsdon et al. 2008). Another technique which is being used to validate annulus formation of long-lived fish species such as the snappers, groupers, and some drum species is the assay of the otolith

core for radiocarbon (^{14}C) resulting from atomic bomb testing in the 1950's and 1960's. This technique is mentioned in Section 4.4.2 of this manual as it relates to accuracy in age determination but without much detail. For more information, Campana (2001) provides an excellent treatise of these techniques and their usefulness in fish age validation.

The intent of this manual is to be a dynamic resource that changes as species specific processing nuances are developed and to serve as a training tool for new laboratory personnel. Descriptions of new and changing techniques will be included in future editions. Standardization of techniques is a cornerstone of fisheries science, and we believe that this manual will facilitate the adoption and incorporation of these techniques and standards for the same and similar species beyond the Gulf region. Moreover, adopting standardized ageing criteria for each species will provide comparable information necessary for age structured stock assessments at state and regional levels.

2.0 Otolith Structure and Function

Most lower vertebrates utilize inner ear elements to process sensory information regarding movement, momentum, spatial orientation, and sound. The dorsal portion of the teleost inner ear includes three semicircular canals each with their own ampulla, a fluid filled chamber for sensing inertia (Figure 2.1A and B). The canals are oriented in such a way as to include the horizontal, lateral, and vertical planes allowing detection of pitch (head up or down), roll (rotation on the head-tail axis), and yaw (head side to side). Movement of the fluid (endolymph) within the ampullae impinges on sensory hair cells lining the walls of the chamber allowing the sensory system to process directional acceleration and deceleration. The dorsal portion also includes the utriculus and the utricular otolith, or lapillus, which is used predominantly to detect gravitational force and sound (Popper and Lu 2000).

The ventral portion of the teleost inner ear includes the sacculus and lagena that each contain their own otoliths, the sagitta and the asteriscus, respectively. This area of the

inner ear appears to be used for both sound detection and acoustic transduction. Sound vibrations differentially affect the otoliths that have a higher density than the fluid filled chambers they occupy. As sound waves are intercepted, the otoliths move independently of the surrounding chamber causing mechanical stimulation of the hair cells (Figure 2.2). This process results in an auditory signal allowing the fish to ‘hear.’




Figure 2.2 Generalized structure and components of the sacculus.

The sagittae are typically the largest otoliths in most fishes and are therefore the most often used for ageing. Please note, however, that some researchers strongly recommend the use




Figure 2.1. A). Location of the otolith pairs within a generalized fish (modified from Secor et al. 1991) and B). medial view of the inner ear (modified from Moyle and Cech 1988).

of other otolith pairs (Secor et al. 1991).

The sagittae lie within the saccula and are attached to a noncellular, olithic membrane. Along the medial surface of the otolith lies a gelatinous pad known as the sulcus acousticus and the nervous tissue called the macula acoustics. This nervous tissue extends from the auditory nerve. Innervation of the gelatinous pad functions to receive stimuli due to angular accelerations, gravity, and sound. Surface features that can be distinguished on some sagittal otoliths include the rostrum and the anterostrum on the anterior end of the otolith and the sulcus acousticus that forms a furrow (sulcal groove) along the medial surface of the otolith (Figure 2.3). The sulcus acousticus can be divided into an anterior ostium section and a posterior cauda section. In some otoliths (e.g., those of certain sciaenid species) a marginal groove is present near the dorsal side of the medial surface of the sagitta.




Figure 2.3. Photomicrograph of medial surface of the right sagittal otolith from a yellowedge grouper (*Epinephelus flavolimbatus*).

Otoliths are crystalline in nature and are built up and outward around a primordium/core region outward by the process of biomineralization, where calcium carbonate, mainly in the form of aragonite, is precipitated on a protein matrix of otolin. The otolin layers are generally oriented parallel to the outer surface of the otolith and are most densely aligned during periods of slower growth

(usually associated with cooler months), thus forming characteristic, concentric **opaque rings** in otolith cross sections (Blacker 1974). Layers that are less densely spaced during periods of faster growth during warmer months make up the **translucent ring** (Figure 2.4). An annual zone consists of one opaque and one translucent ring. When the formation of successive opaque and translucent rings occurs on an annual basis, they are collectively referred to as annual growth zones, and the opaque rings are frequently called the annuli (singular: annulus). Otolith growth in the linear dimension is usually greatest on the axis facing the sagittal midline of the fish.



Figure 2.4. Close up of alternating opaque (O) and translucent (T) rings in a sectioned black drum sagittal otolith under reflected light.

When the alternating rings of an otolith cross section are viewed under magnification, the opaque rings lying along a ‘reading’ or ‘counting’ axis are conventionally the ones tallied for age estimates. The counting axis is generally described by a hypothetical line on one side or the other of the sulcus extending from the core to the outer edge of the otolith (Figure 2.5). Such counting of presumed annuli for the purpose of assigning age estimates is analogous to the practice of dendrochronology, the ageing of trees using tree ring counts from a cross section of the trunk.




Figure 2.5. Transverse section of a black drum sagittal otolith including location of the core and rings along the sulcus. Red dots denote the annuli along the counting axis.

Daily growth increments in sagittal otoliths (first described by Pannella (1971) and later by Pannella (1974), Brothers et al. (1976), Brothers (1984), Campana and Neilson (1985), and Radtke (1989)), are used to infer age, growth events during the first year of life and during specific intervals later in the fish's life. Lapilli have also been shown to provide daily growth increments or rings (Wenner et al. 1990). The astericii are not typically used for daily growth, because they are formed later in life than the other two pairs of otoliths that are present in the fish at hatching/birth.

Otolith morphology differs by species (Figure 2.6). Otolith shape analyses use information extracted from digitized images for species identification (by matching archived key shape descriptors) and, in some cases, to resolve fish populations for the purpose of stock discrimination (Castonguay et al. 1991, Campana and Casselman 1993, Friedland and Reddin 1994, Colura and King 1995, Stransky 2001). The relative size of the otolith also varies widely, but is somewhat based on the needs of the particular species. Pelagic fish which live offshore in clear water tend to have very small otoliths and large eyes relying more on vision than the sensory information derived from the 'inner ear.' In contrast, the nearshore species, which live in much more turbid water, have larger, thicker otoliths since they require more auditory information when sight is limited. Figure 2.7 provides a relative scale of a few pelagic species and their otoliths compared to three species of the drum family, which inhabit the nearshore environments.

In summary, otoliths are anatomical structures that accrete recognizable layers as the result of differential deposition of organic and inorganic material. These layers may correlate with fish growth that varies



Figure 2.6. Variation in sagittal otolith size and shape by species. From left to right: black drum (*Pogonias cromis*), red drum (*Sciaenops ocellatus*), spotted seatrout (*Cynoscion nebulosus*), red grouper (*Epinephalus morio*), sheepshead (*Archosargus probatocephalus*), striped mullet (*Mugil cephalus*), and southern flounder (*Paralichthys lethostigma*).

with time and season and may provide a cumulative historical record of changes in climate, nutrition, hydrographic environment, and other ecological parameters. Their value, to fishery scientists, are as biological and ecological information storage units (akin to “CD-ROMs of fish biology”) that record the temporal signatures of various environmental conditions to which a fish has been subjected from hatching to time of death (Radtke 1990, Kingsmill 1993, C. Wilson personal communication). When comparing otoliths to other fish hard parts; such as vertebrae, scales, finrays, and spines, otoliths often provide more

accurate ageing data due to their continuous accretion and limited resorption whereas other hard parts tend to underestimate age.

The successful application of techniques to enhance the detection of age marks in the otoliths of finfish species is of vital importance in estimating growth and mortality rates, population age structure, and other parameters needed for understanding the population dynamics of important Gulf of Mexico fish stocks and their response to natural phenomena and exploitation.




Figure 2.7 Relative otolith size and body size of several species of Gulf of Mexico fishes. From top to bottom: blue marlin (*Makaira nigricans*), yellowfin tuna (*Thunnus albacares*), wahoo (*Acanthocybium solandri*), red drum (*Sciaenops ocellatus*), spotted seatrout (*Cynoscion nebulosus*), Atlantic croaker (*Micropogonias undulatus*).

3.0 General Processing Techniques

3.1 Otolith Removal

Age data alone is not generally useful to fishery managers unless accompanied by some morphometric, meristic, or other descriptive feature about that fish. Some of these features include: length, weight, sex, and reproductive condition. Otoliths should be removed (post-mortem) after these data are recorded since the otolith removal process will often physically alter the fish, making some of these features impossible to accurately assess.

Sagittal otoliths (the otoliths most commonly used in the Gulf region for ageing) lie inside the otic capsule located toward the posterior end of the ventral surface of the skull. Several methods may be employed to extract otoliths and depend on fish size, shape, and whether or not the whole fish is to be displayed in a market. Some of the more common techniques are described here, as well as in each species account in Section 5.0 of this manual.

In the first method (Figure 3.1A), useful for small fish or when the external appearance of the whole fish must be maintained, the otolith can be excised by cutting into the dorsal junction between the operculum and the body to allow the operculum to be flared open exposing the gills and gill arches (Figure 3.1B). The dorsal attachment of the gill arches and associated tissues to the skull are then cut and the gills and their arches flared forward to expose the tissue surrounding the base of the skull. Under this muscular tissue and lateral to the midline is the outer wall of the otic capsule (Figure 3.1C). Its location and shape varies by species and is described in greater detail in Section 5.0.

Using a stout knife or chisel (depending on

the thickness of the capsule wall), remove layers of the otic capsule wall until the sagitta with its surrounding membrane are fully exposed (Figure 3.2B and C). Use appropriately sized forceps to gently remove the sagitta (Figure 3.2D). Both sagittae can often be extracted through the single opening in the otic capsule. If not, simply repeat the process on the opposite side. If the external appearance of the fish is not a consideration, the gills and gill arches can be removed to expose the otic capsule. The otic capsule can then be scored transversely near its center and broken open along the score to reveal the otoliths.




Figure 3.1. Otolith removal through the gill arches under the operculum; ventral view.

The second method, useful for larger fishes or when the external appearance must not be




Figure 3.2. Removal of the otolith by exposing the otic capsule through the gill cavity using a sharp chisel. A.) Gill cover flared with gills removed exposing otic capsule. B.) Utilization of a chisel or other sharp object to scrape or shave off capsule surface. C.) Open otic capsule with otolith exposed. D.) Otolith removal.

maintained in marketable condition, involves sawing through the dorsal surface of the head, down into or just above the otic capsule (Figure 3.3 Line A). Care must be taken in this method not to shatter the otolith or cut too deep during the initial incision. A hacksaw, heavy knife, bonesaw, or meatsaw is then used to make a transverse cut (Figure 3.3 Line B) from the dorsal side of the head starting just anterior of




Figure 3.3. Cutting planes A. and B. for excision of the sagittal otolith through the upper neurocranium.

where the operculum joins the body (roughly directly above the posterior edge of the preopercular margin). The cut is made deep enough to reach the otic capsule. If the left and right dorsal junctions where the operculum and body meet are cut sufficiently deep, the head can be flexed as if hinged near the snout, exposing the braincase and otic capsule. The otoliths are then removed using forceps.

A third method is the butterfly technique, which is useful on small and medium-sized fishes. This method requires a vertical cut parallel to the long axis of the fish's body (Figure 3.4A). A sharp knife is inserted into the top of the body behind the head and the entire neurocranium is split from posterior to anterior. Once the head is pried opened,

exposing the split otic capsule, the otoliths are removed using forceps (Figure 3.4B). **Note: it is important to make the cut down the center of the head to prevent damaging the otoliths.**




Figure 3.4. The butterfly method.

3.2 Cleaning and Storage

Otoliths have been traditionally used for ageing fish; however, analysis of otolith microchemistry has recently become widespread in fisheries ecology. In order for archived otoliths to be useful for both ageing and microchemistry studies, it is essential that otoliths be properly cleaned and stored to prevent alteration of their chemical composition.

Following extraction, otoliths should be cleaned of any remaining tissue or fluids with water (distilled is preferred). Bleach should not be used because it will dissolve the aragonite matrix and may alter an otolith's chemical composition. Likewise, alcohol should not be used to rinse or store otoliths because it contains trace elements that may penetrate the aragonite matrix of the otolith.

Once cleaned, otoliths should be air-dried completely before storage. Accurate weights (e.g., nearest 0.1 mg) may be determined using an analytical balance. Both left and right otoliths should be stored together in properly labeled paper envelopes or glass/plastic vials and archived for later use. Care should be taken when storing fragile otoliths in paper envelopes. **Note: Storage of specimens in formalin will degrade otoliths by reacting with the protein matrix and should be avoided.** Although left and right otoliths are collected, it is generally agreed that only one side is typically sectioned for ageing. Alternating between left and right for a species could lead to inconsistencies in the ageing process. A comparative analysis between left and right otoliths is recommended for each species since at times the non-designated otolith may need to be used, and there may be a lack of agreement between the left and right otoliths.

Archived otoliths must be assigned unique identification numbers consisting of a species code, a code for the sampling area, and a unique serial number for each individual from the sampling area. This identification number can also include a unique code for the date of capture. In addition, the following information and morphometric data must be recorded for each fish: collection date; location; source (fishery-independent, roving creel, fish house); gear type; length (total, standard, or fork); weight (total or gutted); and sex.

3.3 Sectioning Preparation

The techniques chosen for sectioning otoliths will depend on individual laboratory preferences, budgets, available equipment, and otolith morphology. Three methods of preparation for sectioning are currently used in the Gulf region: embedding whole otoliths in an epoxy resin, mounting a whole otolith to a glass slide, and free hand cutting of whole

otoliths followed by mounting on a slide for sectioning.

3.3.1 Embedding Otoliths

Embedding media are ideal for small or fragile otoliths; however, vapors from these compounds are a potential health hazard so proper lab safety techniques should be followed. Resin mixing, pouring, and processing should be conducted under a fume hood or while wearing a respirator in a well-ventilated area. All individuals exposed to these products should read and have the materials safety data sheets (MSDS) available. Several embedding media are available and are widely used throughout the Gulf States. The most common, Araldite, will be generally discussed, although Loctite (requires UV light to cure) has also been used in a few states for embedding large and small otoliths (Figure 3.5).



Figure 3.5. Small otolith embedded in a block of resin or embedding media that has been removed from the flexible, reusable bullet mold.

3.3.1.1 Embedding Whole Otoliths

Araldite, the more commonly used embedding media, is non-carcinogenic and requires less time to combine the components

for use than some of the resins used in the past in the Gulf region like Spurr (which is discouraged now). To ensure the correct 5:1 ratio of Araldite resin (Araldite-D-US) and hardener (Hardener HY 956 EN/US), stir the contents of each in separate containers and then combine. If only a small amount of epoxy is needed (for a couple of otoliths), resin and hardener should be mixed in a disposable plastic beaker at a 5:1 ratio by weight. Araldite should be prepared under a fume hood or in a well ventilated area while wearing respiratory protection. Avoid skin and eye contact with the resin, hardener, and uncured mixture. As with any potentially hazardous chemical, MSDS should be reviewed and posted in a place accessible to all users.

Araldite should be poured into molds in two steps: a small amount is initially poured into a mold to create a false bottom and left to harden for a day. Next, the sample number is written on the false bottom. An alternative method is to use a permanent ink marking pen to label the inside of each mold with the unique otolith identification number (Figure 3.6). Once labeled, the otolith is placed in the mold, on the false bottom, and covered with a second pour of Araldite. After all the molds on a tray are filled, reposition each otolith as required (correct position is longitudinal; centered with the long axis of the otolith parallel to the sides of the mold and approximately $\frac{3}{4}$ of the way from the square end of each mold) and roll them from side to side to release trapped air bubbles.

Otoliths embedded in Araldite should stand for one hour to allow the reaction heat to dissipate and then be placed in an incubator at 37°C for at least 16 hours while the resin cures. After the resin has completely cured, the otolith blocks are removed from the molds. If a label was applied to the mold or written on the false bottom, it should transfer to the resin

and the blocks do not need to be relabeled. If sample numbers were written on the outside of the embedding mold, this number must be written on the block before it is removed from the mold.



Figure 3.6. Embedding molds labeled with identification information.

3.3.1.2 Embedding Small Otoliths in Bullet Molds

Bullet molds are recommended for small, fragile otoliths (Figure 3.7). Epoxy should be mixed as described above and then added as a thin layer into each cell of the mold with a small metal spatula before the otolith is introduced. The layer of resin on the bottom ensures that the entire otolith is covered and




Figure 3.7 Bullet mold for embedding small or fragile otoliths.

helps to prevent chipping or breaking during sectioning. The otolith is then placed into the cell, centered with the long axis of the otolith parallel to the sides of the cell approximately $\frac{3}{4}$ of the way from the square end of each mold.

This placement ensures adequate material for mounting the block into the saw's chuck. Since the resin is still tacky when the otolith is placed into the mold, you can give the otolith a slight push to fix it into place so the otolith does not move when the remaining resin is added. The blocks should be completely cured as described above before attempting to section.

3.3.1.3 Marking the Core

Regardless of the embedding media or mold type used, marking the otolith core on the resin block is essential for ensuring a transverse cut through the center of the otolith. After a block is removed from a mold, place it under a dissecting scope to locate the otolith core. Though embedded, the otolith should be clearly visible. With an ultra fine point pen or pencil, place a mark over the core of the otolith (Figure 3.8). On one side of the mark, a reference line can be drawn in the transverse plane of the otolith to assist in aligning the blade for sectioning.




Figure 3.8. Embedded otolith with core region marked.

Occasionally, the embedding medium will adhere to the sides of the mold and the block will not be flat on the top side due to the capillary action of the medium. These raised areas can be flattened by sanding them with a small, 1-inch wide belt sander using 100 grit sanding belt, or hand sanding if desired.

3.3.2 Mounting Whole Otoliths on Slides

The following technique works well for both fragile and robust otoliths, but fragile otoliths should be embedded first to prevent breakage. Otoliths to be sectioned should be clean and dry. Prior to sectioning, two slides are made for each otolith. One slide is frosted or has a label applied which contains pertinent information such as species, collection number, and sample number on which the otolith sections will be permanently mounted. The second slide is a plain glass slide that holds the entire otolith during sectioning and is eventually discarded. It is generally necessary to mark each plain slide only with the sample number. As an alternative to glass slides, otoliths can be mounted/glued to heavy cardstock and clipped onto the chuck with minor modifications for production sectioning (See Section 3.4.2.2.1; Figure 3.19).

A whole otolith should be adhered to the plain slide only at the two ends (Figure 3.9). Many labs have used thermoplastic cement. To begin, place the plain slide on a hot plate set at medium to high heat. Apply a small amount of thermoplastic directly onto the slide and allow it to melt. Keep in mind, the slide will have to fit into the saw chuck so it is necessary to leave adequate space at one end of the slide. Remove the slide from the hot plate and be prepared to work quickly, as the thermoplastic will remain malleable for only a few seconds. Scrape the melted thermoplastic into a small pile toward one end of the heated slide using a broad flat instrument. While the thermoplastic

is still soft, place the posterior end of the otolith into the pile of thermoplastic on the slide and pack some over the end of the otolith. If it cools before this can be done, simply return it to the hot plate for a few seconds and then pack. Next, turn the slide around and return the slide to the hot plate being careful not to melt the adhesive just packed on the opposite end. Repeat the above steps while packing thermoplastic around the anterior portion of the otolith. **Remember to leave the core region free of plastic, as this is the area from where the sections will be cut.** Do not try to save time by making a single pile of thermoplastic and splitting it into two smaller piles. This will only make things more difficult later, because the core region may become adhered to the slide as well. This can be especially troublesome with smaller otoliths. An alternative process used by some to adhere the ends of the otolith is to use two-part epoxy or even hot-melt glue applied with a glue gun.

When finished, the otolith should be securely fastened to the slide leaving enough room to place the slide in the saw chuck and ample room to cut sections from the core of the otolith (Figure 3.9).



Figure 3.9. Otolith mounted to a glass slide using thermoplastic on each end. The central portion of the otolith must remain clear of adhesive.

3.3.2.1 Marking the Core

As each otolith is mounted, a line just anterior to the core can be drawn on the otolith

in the transverse plane using a pencil. The line is made slightly posterior to the junction of the ostium and sulcus and is used as a guide during sectioning. Experience will show where to place the reference mark for a given species. An alignment mark may not be necessary on small otoliths, which will have the majority of midsection removed during sectioning.

3.3.3 Thin Sectioning Machine Preparation

Because this saw requires that the otolith be cut free-hand prior to mounting, it is described in greater detail in Section 3.4.3. No pre-sectioning preparation really exists. For most species, it is helpful to make a mark with a pencil anterior to the core to aid in processing on the thin sectioning machine (Section 3.4.3).

3.4 Sectioning Techniques

Otoliths are sectioned typically using rock and gem cutting (lapidary and metallurgical) saws. Three saw types are currently used throughout the Gulf States: the high speed wafering saw; the low speed wafering saw; and the thin sectioning machine. With the wafering saws, thin circular saw blades coated with diamond particles are passed through the otolith in serial cuts to achieve thin sections, which allow the transmittance of light. A tray located directly under the blade is filled with coolant solution. These solutions may be supplied by the saw manufacturer. Alternative saw lubricants include: baby oil, mineral oil, glycerin/water solution, water with a surfactant added, or water only. The thin sectioning machine relies on a larger, single blade to make an initial cut and then the remaining half of the otolith or resin block is adhered to a slide and ground on a second portion of the machine to a single thin section ready to read (Section 3.4.3).

The saws and blades should always

be checked prior to turning them on. It is important to make sure the blade is free of any imperfections that will interfere with sectioning or ruin the blade. After repeated use, each blade should be dressed according to the manufacturer's directions to expose the cutting surface of the diamond particles. A Dremel tool equipped with a fine wire wheel can also be used to clean the flat portion of the blade.

Never start the saw with the resin block resting on the blade. Allow the saw blade to achieve target speed before making contact with the sample. Failure to do so could result in a broken blade or in the case of whole mounted otoliths, stripping the sample from the slide. Make sure to read all directions provided by the saw's manufacturer.

3.4.1 High Speed Wafering Saw

A high speed saw (Figure 3.10) has several advantages in terms of production; however, it is one of the more expensive saws, and blades are costly (see Appendix 8.2). Problems with electrolysis or corrosion between the aluminum saw blade flanges and the copper-coated saw blades have been encountered, but do not



Figure 3.10. High speed wafering saw (cover opened).

appear to impact saw operation or blade life. Saw blade flanges may have to be replaced every three years.

A high speed wafering saw with a maximum speed of 5000-rpm (in 100-rpm increments) can be used with four- or six-inch, diamond-coated saw blades to produce a section thickness of 0.5 mm. The high speed saw has a 1000-g load capacity (in 10-g increments) and allows for chuck crossfeed adjustments in 0.005-mm increments. The saw blade is lubricated and material residue is flushed away by a recirculating lubricant stream from a submersible pump (Figure 3.11). Loss of lubricant due to spray off the high speed blade is prevented by a cover. The cover also shields the operator from the high-speed blade, airborne material particles, and lubricant aerosols from the cutting operation. Sample sections are retained in a metal basket over the lubricant reservoir but may be difficult to locate as they sometimes spray off the blade and adhere to the interior surface of the cover.

Sectioning times through a resin-embedded otolith on the high speed saw will vary based on block size, but usually take from 15 to 45 seconds. Cutting speed, load, and chuck position are controlled by pressure pads and settings for all three are displayed digitally and will need to be adjusted for each species being processed. One downside to the high speed wafering saw is that the saw has a safety switch which prevents blade or pump operation when the cover is open and eliminates the ability to manually section otoliths or dress the blade.

3.4.1.1 Embedded Otoliths

The resin block containing the otolith is placed in the chuck of a high speed wafering saw equipped with either a four- or six-inch diamond blade. The block is oriented so the long axis of the otolith is perpendicular to the



Figure 3.11. High speed wafering saw showing blade, cutting arm, coolant reservoir, and pumps with resin block ready to be sectioned.

saw blade and the anterior end of the otolith is nearest the chuck (Figure 3.11). Sectioning begins just posterior to the otolith core, and sequential sections are made approaching the core region until a good section is obtained. The otolith block is advanced approximately 1 mm toward the saw blade after each cut which produces about a 0.5 mm thick section. Sectioning is typically done at 3000 rpm with a 1000-g load, and typically takes less than 30 seconds for all otolith sizes. Sections are examined under a dissecting microscope to identify the section containing the otolith core, which is then affixed to a labeled glass slide (see Section 3.5).

3.4.1.2 Whole Otoliths

Whole mounted otoliths can not be cut with this saw since the cover must remain closed during sectioning.

3.4.2 Low Speed Wafering Saw

There are several benefits associated with low speed wafering saws. Simultaneous operation of several of these sturdy saws enables a high sectioning production rate. Low speed wafering saws are less expensive than the higher speed models. They are relatively safe, require no safety shield, are simple to operate, and relatively quiet. The low speed saws have a maximum speed of 300 rpm and generally use four-inch, diamond wafering blades with a 0.3-mm kerf.

Most of the saw manufacturers provide repair services and technical support and will recommend appropriate sized chucks for various sizes of resin blocks. Finally, the small size of these units allows for transfer between laboratories. Disadvantages are that the diamond wafering blades are fragile (brittle), expensive to replace, susceptible to bending and chipping, and processing time can be relatively long for extremely large or embedded otoliths. A number of laboratories have begun utilizing multiple blades with spacers on the same low speed saw to obtain simultaneously cut sections for 'production sectioning' (Section 3.4.2.2.1).

3.4.2.1 Embedded Otoliths

A resin block containing a single otolith is positioned in the saw chuck so the cut will result in a 90° cross sectioning of the otolith very near, if not through, the core. The chuck may be adjusted to orient the block by loosening the Phillips head screws (or thumbscrew if the saw has a vise-type sample holder) on the specimen arm. The operator should view the block from the top or bottom as well as from the front to check for alignment. When the block is correctly aligned, the screws are tightened (Figure 3.12). Failure to tighten the block in the chuck appropriately may result in a ruined blade. Every effort should be exercised during

preparation to have the otolith properly aligned in the block to avoid having to make substantial adjustments to achieve the correct orientation in the saw chuck. For otoliths embedded in small bullet molds, it may be necessary to first mount the block onto a slide using thermoplastic or other adhesive and then align the slide in the chuck.

Sectioning begins posterior to the otolith core near the junction of the ostium and sulcus and sequential sections are made approaching the core until a good core section is obtained (Figure 3.13). The block is moved across the blade after each cut using the micrometer cross feed to adjust the desired thickness of each



Figure 3.12. Embedded otolith mounted in low speed saw.

section (Figure 3.14). Depending on the type, size or fragility of the otolith and embedding medium used, the saw speed can be adjusted using the speed control, and weight may be added or removed from the specimen arm to achieve the best cut. With practice, a section



Figure 3.13 Resin block with otolith oriented against a blade at 90°.

containing the core region should be reached within two to three cuts.

Sections are removed from the specimen tray, rinsed in water, dried, and viewed under a low-power microscope to verify that a good core section has been obtained. If the core was missed, the block may be rechucked in the saw to attempt another section.




Figure 3.14. Adjusting the alignment of the block with the micrometer.

Permanently affix the best section or sections to the final, labeled slide using an appropriate mounting medium and set aside to dry or cure (reference Section 3.5) (Figure 3.15).

3.4.2.2 Whole Mounted Otoliths

When sectioning whole mounted otoliths (Section 3.3.2) using a low speed wafering saw, check the recommended arm weight and blade




Figure 3.15. Mounting of otolith core section on final slide.

speed for that species (some specifics are provided in Section 5.0). This may require some trial and error with new species. Secure the slide with the adhered otolith in the chuck with an Allen wrench so that it does not slip during sectioning, but do not over tighten as the slide can break. Also check the angle of the chuck to ensure that the blade will section the otolith in the transverse plane. Line up the blade based on the core which should have had its position marked with a pencil prior to mounting. Although it is not necessary, it is often

easier to begin sectioning with the micrometer crossfeed scale at the zero position.

To begin sectioning, turn the saw on with the otolith raised above the blade (specimen arm in the up position). Do not start the saw while the otolith is resting on the blade as it could damage the otolith and/or the blade. Gently lower the otolith onto the turning blade and begin sectioning. Depending on the species, size of the otolith, weight, and saw speed, it can take anywhere from 30 seconds to several minutes to cut through the otolith (Figure 3.16).



Figure 3.16. First transverse cut of a whole mounted otolith.

It is practical to cut three or four sections from the otolith to ensure a section that includes the core. When the blade passes through the otolith and begins to cut the glass slide, lift the specimen arm off the blade and advance the saw blade through the core (Figure 3.17).

Sections are generally cut in 0.5-mm (500- μ m) increments. However, this thickness can be altered depending on the species (see Section 5.0 for specific recommendations).

Once all sections have been cut, lower the specimen tray and rotate it out from under the blade. Pull the specimen basket out of the



Figure 3.17. Serial cuts from a whole mounted otolith.

cutting solution and remove all otolith sections with forceps. Rinse the sections in water and allow them to dry. Examine the sections under a low-power microscope to ensure that a good core section has been obtained. Affix the section or sections to the final slide using an appropriate mounting medium (Figure 3.18). It is best to position otolith sections on the final slide in a consistent manner for easier ageing. It may be beneficial to affix the sections in such a way that they are not permanent (see Section 3.5 for options and benefits).




Figure 3.18. Three otolith sections prepared for the final mount and ring enumeration.

3.4.2.2.1 Production Sectioning

A number of laboratories have begun utilizing multiple blades with spacers on the same low speed saw to obtain simultaneously cut sections from a single pass. This technique works very well for both fragile and robust otoliths. As noted previously (Section 3.3.1.2), fragile otoliths should still be embedded before sectioning to prevent breakage. The advantage of this multi-blade technique is that it results in three or four sections which should contain the core or at least be very close to the core in one-third to one-quarter of the total processing time.

To section the otolith with multiple blades, whole specimens are hot-melt glued to a blank slide or small pieces of tag paper or card stock cut to the size of a standard slide (Figure 3.19). If using a blank slide, use the same technique as noted in Section 3.3.2 with the existing chuck. Paper slides are held to the saw arm by a small binder clip (Figure 3.20). In both cases, the specimen is slowly lowered onto the spinning blades of the saw with the blades running through the water bath for lubrication.



Figure 3.19 Whole otolith ‘glued’ to cardstock for production sectioning.

Thin, transverse sections are cut with a low speed saw. Generally, 3 to 4 blades, each separated by a 400- μm spacer, are used to yield 2 to 3 transverse sections approximately 300-400 μm in thickness (Figure 3.21). The sections are then recovered from the basket, dried, and affixed to a final, labeled slide using a mounting medium (Section 3.5). Spacers may be difficult to locate for purchase but can be made easily in the lab by cutting the center sections out of old blades. The spacer allows the blades to run simultaneously and cut consistent sections that often don’t require additional sanding.




Figure 3.20 Binder clip holding cardstock in chuck preparing to be sectioned by multiple blades.

3.4.3 Thin Sectioning Machine

The thin sectioning machine is primarily used to section unembedded, whole otoliths. The procedure borrows petrographic techniques from geology and reduces sectioning time by eliminating the time-consuming steps of embedding and polishing (See section 3.7.1). In addition, the apparatus allows the technician




Figure 3.21 Four blades set up with spacers for production sectioning.

to prepare a large number of otoliths at one time. The thin sectioning machine can be used to create ‘frosted’ slides by grinding one end of a less expensive, clear blank slide on the machine’s lap arm pad. **Note: The sectioning process is quite loud so ear plugs or other protective ear wear is recommended.**

The water-cooled, thin sectioning machine is equipped with two individual tools; a cut-off saw and a precision grinder (Figure 3.22). The saw is equipped with a 20-cm diamond blade while the grinder is equipped with a 20-cm, vertically mounted, 320-mesh, metal-bonded-diamond grinding lap. The grinding lap is fitted with a precision dial controlled thickness gauge allowing the technician to vary the section thickness. Both have aluminum guide arms for feeding slides to the blades.

The following is a method for the rapid processing of large otoliths first described by Cowan et al. (1995) with some minor modifications. Otoliths are hand held and cut along the transverse plane near the core using the cut-off saw before mounting onto slides (Figure 3.23). To ensure a high quality section, it is imperative to cut as close to the core



Figure 3.22. Thin section machine containing a high-concentration-diamond, continuous-rim-blade cut-off saw (left) and a precision grinder (right).

as possible without actually cutting through it so that the core is contained at the transverse plane edge of the otolith half to be mounted. The cut surface of the otolith half is then pressed against the precision grinder to remove any rough edges or scratches. Additional polishing may further reduce scratches. This will provide a readable surface on both sides of the finished section.



Figure 3.23. Hand cutting an otolith on the high speed thin sectioning saw.

Allow the otolith half containing the core to dry and mount it cut side down onto a final

microscope slide. For ease of processing, two otoliths can be mounted per slide with identification numbers written under each using a water-proof marker (Figure 3.24).

After curing, the slide containing the otolith halves is placed in the guide arm of the cut-off saw and guided past the saw to remove all but approximately a 100 μm section of each of the otolith halves. The slide is then placed into the precision grinder guide arm and fed past the grinding lap to remove any rough edges or scratches (Figure 3.25). Once the slides are dry, the otolith sections on each slide may be covered with a few drops of mounting medium which may eliminate the need for polishing. The otoliths are then ready to be read.



Figure 3.24. Otolith halves mounted on microscope slides with Loctite which is cured under ultraviolet (UV) light.

The following technique can be used for fragile (e.g., flounder) or small otoliths (e.g., mullet) and is similar to processing larger otoliths, but requires greater manual dexterity as all processing is done on the precision grinder. Marking the core is essential in achieving a quality section using this technique. Otoliths are handheld by the posterior end and ground down to the transverse plane near the core. Again, it is imperative to get as close to the core as possible. The otolith half is mounted

cut side down onto a labeled microscope slide and cured. After curing, the slide is handheld and pushed against the grinder until remaining material is removed to approximately 1 cm. The slide is then placed into the precision grinder guide arm and fed past the grinding lap to reduce the section down to the desired thickness.



Figure 3.25. Final polishing of otolith sections using grinding arm.

3.5 Mounting Sections

Section mounting, or adhering the otolith sections to a glass slide, can be done in several ways. The two most common mounting media used in the Gulf region are thermoplastic cement and Flo-Texx, although several labs are moving away from thermoplastic. Hand or machine polishing to remove saw abrasions or other imperfections from the section surface can take place before and/or after mounting. Following mounting, it is useful to apply a coat of commercially available permanent coverings such as Flo-Texx or temporary coverings such as immersion oil, glycerin/water solution, or plain water to increase clarity when reading. **Note: The use of Histomount for slide preparation is strongly discouraged due**

to its tendency to discolor and crack over time.

To temporarily mount sections, thermoplastic works well as long as the section is not covered with any kind of permanent covering agent. In the event one would want to release or move the section, the thermoplastic can be melted again and the section adjusted, removed, or replaced on the slide.

3.6 Alternative Techniques

3.6.1 Whole Otoliths

Examination of a whole otolith using transmitted light can often reveal marks expressed on the surface (Figure 3.26). This technique has predominantly been used for otoliths taken from larval and small fish, but has been used successfully to age older gag (*Mycteroperca microlepis*; McErlean 1963) and red grouper (*Epinephelus morio*; Johnson and Collins 1994). In general, marks observed from whole otoliths may correspond with opaque rings observed from sectioned otoliths, but this is not always the case. The use of



Figure 3.26. Whole otolith from hardhead catfish, *Arius felis*. Rings are apparent from outer surface.

whole otoliths requires less time and effort than sectioned otoliths, but validation must be undertaken to verify that rings counted on whole otoliths correspond with the ‘correct’ number of rings observed in sections. Rings counted on whole otoliths for striped mullet (*Mugil cephalus*) in Mississippi were consistently one ring fewer than the ring counts from sections of the same otoliths (J. Warren personal communication). Inconsistencies have also been observed when comparing whole and sectioned southern flounder (*Paralichthys lethostigma*) otoliths (A. Fischer personal communication).

Whole otoliths can be read using a dissecting microscope and either reflected or transmitted light. In most cases, 12x magnification is used, but 6x magnification may provide a ‘cleaner’ image. **Note: Switching between the two magnification lenses while reading the whole otolith may produce better results.** Additionally, adjusting the angle of illumination or otolith position may increase the contrast of the rings versus the increment. Otoliths can be placed in a small watch-glass with enough water, oil, or glycerin/water to completely submerge the otolith to enhance the marks. An alternative to using a watch-glass would be to use a plastic tissue culture tray. The advantage to culture tray cells is that multiple otoliths could be viewed simultaneously and the individual walled cells prevent the otoliths from “mixing” if the dish were to be bumped or moved accidentally. When reading whole otoliths, the younger fish are easier to age. The first annulus is generally clearer and whole otoliths from fish beyond age-5 become progressively more difficult to age as one gets further from the core. The ventral, posterior edge of the otolith is usually a better area to read; however, adjusting the angle of the light source or orientation of the otolith may produce better results (Figure 3.27).



Figure 3.27. Ventral posterior edge of a whole sagittal otolith from an age-5 king mackerel.

3.6.2 Break and Burn

As an alternative to thin-sectioning sagittal otoliths, fish ages can be determined using the “Break-and-Burn” method (Christensen 1964). With this method, the sagittal otolith is literally broken in half through its nucleus (core), and the exposed surface is heated over an alcohol flame to enhance the contrast between the organic and inorganic components of the matrix. Manual manipulation of an otolith half using fine-tipped forceps is required so this method is usually limited to larger otoliths (>8-10 mm in length). This does not preclude using this technique on smaller otoliths, but it does require more skill and care in the burning process. This method is successfully being used on white grunt (*Haemulon plumieri*) and red porgy (*Pagrus pagrus*) and may be appropriate for other species when rapid production ageing (i.e., to year class) is required, rather than specific ageing for growth (D. Murie, personal communication).

To break a sagittal otolith, the transverse plane of the otolith is scored through the nucleus using a diamond-edged pen and then snapped in two using finger pressure. The broken surface of one-half of the otolith is then held at an angle and moved back and forth above an alcohol flame. **Note: When burning the surface it is important to keep the flame**

evenly distributed over the otolith’s surface to get an even burn. The otolith should not touch the flame directly or it will burn too quickly and char the surface making ageing impossible. The time required to burn a surface depends on the species and size of an otolith, but is usually no more than 10-15 seconds. Care should be taken with smaller otoliths as they will require less time.

This process differentially burns the organic matrices within the annuli of the otolith, with the translucent bands of slow growth burning dark relative to opaque bands of faster growth (when viewed under reflected light). The otolith half is cooled (usually less than 30 seconds) and pressed into a dark-colored plasticine block (blue or green works well) with the burnt surface upright and tilted slightly (Figure 3.28).



Figure 3.28. A break-and-burn otolith pressed in a plasticine block under a reflected light source.

3.6.2.1 Reading and Interpretation

Bands are counted using a stereomicroscope fitted with a fiber optic light source (reflected

light) positioned to reduce glare while providing a source of focused light when using high power (Figure 3.29). The contrast among the bands can be enhanced using a drop of vegetable or canola oil on the burnt surface.

Age estimates for a series of white grunt otoliths were processed with this method as well as sectioning and were almost 99% in agreement at least up to 16 years of age (Murie and Parkyn 1999). The primary advantage of the break and burn technique over thin-sectioning is the greatly reduced amount of time required for processing otoliths (minutes rather than hours). In addition, long-term storage of burnt otoliths does not appear to result in the fading of bands (D. Murie personal communication). Otoliths can be re-burnt to enhance visibility of bands or in most cases, the other half of the otolith can be used.



Figure 3.29. Broken-and-burnt surface of a sagittal otolith from an age-6+ white grunt under reflected light (N=nucleus and S=sulcus) (from Murie and Parkyn 1999).

3.6.3 Scales

Scales have long been used for ageing fish and are one of the earliest hard parts to be found with well defined rings, which were supposed to be annuli. During the early 1900s the use of scales for ageing fish and separating fish populations led to seminal research in ecology and fisheries management (Sinclair 1988). By the early 1920s, Welsh and Breder (1924) reported age

and growth information for fish from southwest Florida using scales. Age determination using scales was so common that Lee (1920) reviewed their successful use for a variety of species.

3.6.3.1 Scale Collection and Handling

Scales are often removed from the middle of the body, below the dorsal fin (Figure 3.30), but many species have precedent for removing scales from other locations. It is necessary to collect scales from a region of the body where scales first form. When removing scales from live fish, however, the collection area should be 're-slimed' to aid healing the fish's epidermis; it is recommended to use a wet and bare finger to spread the fish mucus back over the area where fish scales are collected. Because some scales are unsuitable for ageing, it is recommended that one collects 6-10 scales per fish. A typical problem arises from regenerated scales that are missing the interpretable ridges (i.e., circuli and radii) that define the annuli in the central portion (i.e., around the focus) of the scale.




Figure 3.30. Six to ten American shad (*Alosa sapidissima*) scales are taken from an area below the center of the dorsal fin and above the lateral line.

3.6.3.1.1 Raw Scales

One of the advantages of using scales in favor of other anatomical parts is that samples can be obtained without affecting the appearance of a fish in the market or sacrificing the fish in the field. Another advantage of using scales is that they are easily collected and stored. Scales can be removed quickly by using forceps or a knife

and stored in inexpensive envelopes. If long-term storage is anticipated, it is recommended that scales are cleaned and stored in a cool, dry area and that moth balls are included to prevent mites from damaging the scales. Cleaning the scales when initially collected can save time later. A small brush, such as a toothbrush, and a cleaner (e.g., a mild soap solution, alcohol, or diluted bleach) will be necessary to apply to the scale once the fish slime has dried. No further processing is necessary if the raw scales are examined directly, although some additional effort to mount the scale, either dry or wet, or make an acetate impression of the scale can enhance the details of it for viewing and interpretation. Enhancement techniques are described in much greater detail by Dery (1983).

3.6.3.1.2 Scale Impressions

The sculptured side of a fish scale can be imprinted on laminated plastic by using pressure, such as with a roller press (Figure 3.31). Cellulose acetate can also be used, but this medium requires heat, heavy pressure, or softening chemicals for impressions. Making impressions is a more laborious technique, but the time and cost can often be justified and provide several advantages over raw scales. First, impressions can enhance the details of scales with delicate features. Second, the impression will be flat, even if the scale is curved. A flat image reduces problems associated with light diffraction and minimizes the focal depth of field necessary for recording good photographs or digital images. Third, larger scales may be too thick to be transparent enough for direct viewing while impressions can be viewed using transmitted or reflected light. Fourth, multiple scale impressions on a single slide can be easier to handle than many small, loose scales in an envelope, and the best scales can be easily selected for reading. Fifth, impressions can be archived indefinitely.




Figure 3.31 Scales on acetate passing through roller press to make impressions.

Clean and dry scales are typically placed on a blank slide in order and a single acetate slide is laid over them. Carefully, place the sandwiched scales into the rollers of the press (Figure 3.31). As the machine passes them through, the scales are pressed into the acetate and the result is a negative relief of the scales surface features. The acetate is removed from the slide and the scales, which may stick to the acetate, are removed and returned to their archive envelope returned to



Figure 3.32 Final scale impression on acetate slide.

their archive envelope for storage. The acetate slide is labeled and can now be read (Figure 3.32).

3.6.3.2 Scale Reading and Interpretation

Scales or scale impressions can be viewed with a light microscope, a microfiche reader, or a microprojector (Figure 3.33). The generalized criteria for counting annuli are to examine the patterns of cutting over, discontinuity, or crowding of the circuli. **True annuli appear as lines on the scale surface and follow the contour of the scale periphery** (Figure 3.34). They are most evident along the lateral fields of the scale. False annuli are generally faint in comparison to true annuli, but one important exception is the fresh-water zone mark laid down when juvenile shad move from fresh to salt water at the end of their first summer.

The approximate locations of the true annuli during the pre-spawning period of a fish's life can be found by counting the number of transverse grooves that follow the baseline groove (Figure 3.34). The transverse grooves are unreliable for ageing the fish. The anterior edge of each true annulus generally falls between a narrow range of transverse grooves for virgin fish.

Additional rings, such as spawning marks, can be mistaken for annuli when interpreting scales. For example, in the south, American shad (*Alosa sapidissima*) are semelparous, so spawning marks do not appear, but spawning marks need to be identified correctly for iteroparous populations of American shad (i.e., populations north of the Carolinas). Spawning marks are more jagged in appearance than true annuli because they arise from both eroding and regenerating processes of the scale margin. As a fish ages, the space between consecutive annuli becomes narrower, and the erosion caused by spawning can obliterate the recent annulus. Thus, after maturation the spawning mark is the annulus



Figure 3.33 Scale impression being read on a microfiche.

mark in species like American shad that spawn immediately following annulus deposition in its northern range. Spawning marks in American shad do not occur on the otolith, only on the scales, thereby offering a specific advantage over otoliths (i.e., these spawning marks indicate the age and size at maturation and the number of years a fish has spawned).

Several other species in the southeast United States have been successfully aged using scales; Atlantic menhaden (*Brevoortia tyrannus*), ballyhoo (*Hemiramphus brasiliensis*), bluefish (*Pomatomus saltatrix*), dolphin (*Coryphaena hippurus*), tomtate (*Haemulon aurolineatum*), knobbed porgy (*Calamus nodosus*), whitebone porgy (*Calamus leucosteus*), black drum (*Pogonias cromis*), red drum (*Sciaenops ocellatus*), southern kingfish (*Menticirrhus americanus*), and striped mullet (*Mugil cephalus*). In these species, scale annuli were validated using marginal increment analyses from recaptured fish or were judged to be valid based on the appearance of continuous growth of the scale and circuli patterns.




Figure 3.34. An acetate impression of a scale from an age-6 American shad (*Alosa sapidissima*) collected in the York River, Virginia, depicting annuli (Roman numerals), transverse grooves (Arabic numerals), and other features.

Validation of scale annuli is essential because scales may not always be useful for ageing fishes. Beamish and McFarlane (1987) demonstrated that the scale method provided erroneous ages for 16 freshwater and marine species. In general, maximum scale ages underestimated validated ages or ages determined by some alternative method (i.e., otoliths). Otoliths continue to grow as a fish ages. Problems can arise using scale, however, as they do not continuously grow in older fish and the calcium in the scales can be resorbed in stressed fish. Scales are regarded as unsuitable for ageing large pelagic fishes, namely tunas, billfishes, and sharks (Casselman

1983). Lowerre-Barbieri et al. (1994) published a good example of how to compare scale and otolith methods, and they noted that crowding of annuli on the scale margin was problematic in older weakfish (*Cynoscion regalis*). They concluded that sectioned otoliths provided more accurate ages and more precise indications of annulus location. Secor et al. (1995b) concluded that scales were suitable for ageing striped bass (*Morone saxatilis*) younger than 12 years. They noted, however, that most stock assessments for this species are still based on scale ages to avoid sacrificing the oldest and largest females which serve as broodstock. To compensate for the

use of scales instead of otoliths, they reported a linear equation that could correct the ages of older fish. These examples should make it clear that before expending time, energy, and funds to collect and use scales for life history studies or stock assessments, the issue of validating annulus formation on scales should be addressed (Section 4.4.1).

In summary, scales are not appropriate for ageing many species, particularly slow-growing, long-lived species. However, scales may be useful for ageing faster-growing, short-lived fishes, and for ageing younger individuals of slower-growing species when mortality from scientific sampling needs to be reduced or eliminated. Using scales has some advantages over other hard parts, such as their ease to collect, store, and process without sacrificing the fish. Validation of annulus formation is necessary, however, to make use of these advantages of scales for ageing fish. Although otoliths have been demonstrated to be quite reliable for determining age, scales may become more widely used in the future where non-lethal sampling is desirable or required. In addition, scale shape has been used for stock identification for several decades (Ihssen et al. 1981), and recently Moran and Baker (2002) demonstrated that archival scale samples are valuable for genotyping historical collections. The historical use of scales and the familiarity that most fish biologists have with scales have led to archived material at many labs, and these historic and newer collections can continue to play a part in understanding the population dynamics of fishes.

3.6.4 Spines and Finrays

Using spines and finrays for age and growth studies offers certain advantages over otoliths and other hard parts. In most cases, spines and rays can be removed and processed more easily than scales and otoliths. It is rarely necessary

to sacrifice the fish or significantly mutilate the carcass when sampling, which could reduce the market value of a commercially harvested species. Soft finrays are especially useful because they can be removed at the time of tagging and compared with the corresponding structure at the time of recapture. For more detail on these techniques see Casselman (1983). Unlike scales, the annuli in finrays and spines remain prominent for older fish when scale annuli are no longer identifiable.

Although spines and finrays can be useful in the estimation of age and growth in fish, there are disadvantages. In older fish the core can undergo resorption and become vascularized, thus obscuring or eliminating the first few annuli resulting in an underestimation of age (Figure 3.35). This is common in many of the oceanic pelagic species. Spines and finrays from older fish are also similar to scales in that the distal translucent zones may be so close together that they appear to coalesce, making optical resolution difficult or impossible. **Note: Successfully determining age from spines and finrays requires that the structures be sectioned near their base in a precise transverse plane, although the exact location of the section depends on the species.**

False annuli, or pseudoannuli, appear similar to annuli but are associated with checks and zones that are often incomplete and irregular, and frequently found only in one region of the structure. Although they may be prominent, pseudoannuli are not associated with growth zones that form during the principal annual cessation or reduction in growth that produces annuli and should not be counted when ageing. Validation of the hard part for each new species is necessary to confirm that observed marks are, in fact, produced annually (See Section 4.2.1 and Section 5.6).




Figure 3.35. Resorption and deterioration in the core (indicated by dashed lines) of the first dorsal spine of a cobia (*Rachycentron canadum*).

3.6.4.1 Sectioning Spines

While many different methods for sectioning fin spines exist, two techniques have been used successfully in the Gulf states using either the thin sectioning machine (AMRD, LSU) or a low speed wafering saw (FWRI). Differences in method between the two saws are detailed below. The thickness of the transverse section must be adjusted to assure that annuli are visible. Sections may be soaked in solutions containing acetic acid or bleach to remove unwanted tissue from their surface to make annuli observation and quantification easier. Spine and finray sections are then mounted using any one of the techniques mentioned in Section 3.5.

3.6.4.1.1 Thin Sectioning Machine

The shaft of each dorsal spine is sectioned slightly above the condyle. The exact location in each species is determined by trial and error. A section too far up the spine (Figure 3.36, Line A) will result in more closely spaced annuli, and a section made too close to the condyle will result

in annuli that are obscured by the convolutions in the condyle of the spine (Figure 3.36, Line C). **Note: If the section is made below the lateral groove, the annuli will be truncated and it will be impossible to follow them all the way around the spine.**

To make a transverse section, hold the spine horizontally and perpendicular to the saw blade. Then make the first cut while holding the spine as steady as possible (Figure 3.37). Push the spine into the cutting wheel with light pressure (forcing the spine cause fissures/cracks). This will result in two portions of the spine; the distal portion and the proximal portion containing the condyle. Next, polish the cut surface of the distal portion and mount it to a final microscope slide with thermoplastic so that the plane of the cut is parallel to the plane of the slide (Figure 3.38). Place the slide into the guide arm of the cut-off saw. Make the second transverse cut




Figure 3.36 The first transverse cut (line B) provides the most widely spaced annuli with the best resolution when separating the condyle from the shaft (line A will result in more closely spaced annuli and line C will result in annuli which may be obscured by the condyle).



Figure 3.37. Freehand transverse cut of spine on thin sectioning machine.




Figure 3.38. Distal portion of spine cemented to slide, ready for sectioning.

using the guide arm to pass the spine over the blade (Figure 3.39). The result is a spine section approximately 0.5-1.0 mm thick. The thickness can be adjusted by placing the slide in the guide arm of the grinding lap and feeding the section back and forth to polish it. Have a microscope set at 40x magnification nearby to monitor the clarity of the section as you adjust the thickness of it.

3.6.4.1.2 Low Speed Wafering Saw

The second technique for sectioning a spine uses a low speed wafering saw and is similar to the methods described for whole mounted otoliths (Section 3.4.2.2).

The whole spine is attached laterally to a glass slide with thermoplastic at the condyle and the tip, making sure that the shaft itself is free from the slide (Figure 3.40).

The slide is then placed in the chuck of the saw and lowered onto the spinning blade to section the spine (Figure 3.41) in a transverse plane at 0.75 mm intervals beginning at the shaft base (just above the condyle). As many as six serial sections can be mounted on microscope slides (Section 3.5).

3.6.4.2 Sectioning Finrays

Finrays used for age determination are typically removed from the dorsal or pectoral fin. A modification of the method of Chilton and Beamish (1977, 1982) has been used successfully with finrays to estimate ages for white grunt (*Haemulon plumieri*) up to about ten years (Murie and Parkyn 1999).

Soft finrays are removed from the dorsal fin (rays 4-7) by cutting across their base. For live fish, the rays must be removed as close to the dorsal surface of the body as possible to make sure



Figure 3.39. Removal of excess material from mounted spine.



Figure 3.40. Whole dorsal spine mounted to slide.

that all annuli (especially the first) are present in the base of the ray. On dead fish, the rays can be removed down to their base (knuckles), which extends into the muscle of the fish. Finrays should be trimmed of excess tissue and placed in a non-gummed manila coin envelope with the cut surface exposed to the air and the finrays lying parallel to one another to dry for two to five days (Figure 3.42). **Note: It is important to arrange the finrays in a parallel position so that they can be processed without having to be cut apart and realigned.**




Figure 3.41. Whole spine mounted laterally to slide with thermoplastic and positioned for transverse sectioning on a low speed saw.

Once dried, the finrays may be embedded using a two-part epoxy resin (Figure 3.43). Though embedding is necessary to hold the finrays in the saw chuck, the use of a mold is not necessary. The finrays are placed on a piece



Figure 3.42. Dorsal finrays from a white grunt arranged for drying in a coin envelope.

of parafilm (to which resin does not adhere) and resin is applied over the basal surface of the dried finrays. Finally, the distal portion of the finrays should be embedded in a large resin tear-drop. Once cured, four to five serial sections (0.5-0.8 mm) can be cut from the distal end of the finray block. The sections are permanently mounted on a labeled slide for annuli enumeration (Figure 3.44).

3.6.4.3. Reading and Interpretation

As with other hard part sections, the finray sections are best viewed using a compound microscope, although they can be projected with a microfiche projector or viewed using a microscopic video camera and monitor.

As with spines and scales, there is a problem of annuli accumulation at the edge of the finray structure, which can lead to an underestimation of the true age of the fish (Figure 3.45). For example, white grunt age estimates obtained from finrays and sectioned otoliths agreed in 90% of the readings only for fish less than ten years old and decreased to 13% agreement for fish between 11 and 18 years of age. Finrays from these older fish do not display significant growth between consecutive annuli, making it difficult to count individual annuli near the edge of the structure.



Figure 3.43. Embedded dorsal finrays from a white grunt (rays are obscured by cured resin). Sections have been removed from the basal portion (right side) of finray block.

The degree of potential ageing bias due to underestimation should be evaluated for each species as the peculiarities of species-specific growth will affect the observed annuli pattern in the finrays.



Figure 3.44. Sections of dorsal finrays from a white grunt mounted to final slide.

3.6.5 Vertebrae

In fish (i.e., elasmobranchs) that lack hard parts such as otoliths or usable scales, age and growth information is derived from marks observed on vertebral centra and spines (Cailliet 1990). The current hypothesis is that thin, opaque bands are formed in the winter months and broad, translucent bands are formed in the summer months, although this has only been validated for a few species.

3.6.5.1 Extraction and Storage

Approximately ten vertebrae should




Figure 3.45. Magnified cross-section of the dorsal finrays from an age-6 white grunt.

be removed from just below the dorsal fin (Figure 3.46). This is accomplished by cutting straight through the body just anterior and posterior to the dorsal fin. The removed vertebrae can be cleaned of excess tissue and separated using a sharp knife or scalpel blade (Figure 3.47).




Figure 3.46. Depiction of shark with rectangular area denoting the section of vertebrae that should be removed.

The individual discs are soaked in a 5% sodium hypochlorite solution for 5-30 minutes or until all connective tissue has been removed. Once clean, vertebrae are stored in 70% ethanol until individual vertebrae are utilized for ageing.



Figure 3.47. Separating and cleaning vertebrae of excessive tissue before sectioning.

3.6.5.2 Sectioning and Reading

Depending on the species of fish, vertebrae can be cut in half (perpendicular to the centrum face), read whole, or cut into thin sections. When cut into thin sections, the vertebrae may be stained to enhance the contrast of growth bands. The sections are then mounted to a slide and examined with a dissecting microscope (Figure 3.48).

Most coastal sharks are born in spring to early summer. When ageing sharks, one must remember that the first band observed in a vertebral section is called the birthmark and is theoretically formed at the time the shark is born (Figure 3.48). The second band is formed six months later during winter, and a new band is formed every winter following. Therefore, a shark with two opaque bands is approximately 6+ months old, but is still considered an age-0.

3.7 Section Enhancement

When reading otolith, finray, spine, or vertebrae sections, saw marks and other surface scratches can often reduce the reader's ability to see rings clearly. Optional techniques to enhance the readability of otolith sections include polishing, etching, staining, clearing,




Figure 3.48. Sectioned vertebrae of an 8.5 year old shark with (birthmark indicated).

and baking. Other enhancement techniques may improve readability without directly affecting the otolith section such as alternative lighting types, filters, polarizers, and light sources. The resolution on most otoliths can be improved using one or several of these techniques; however, a bit of trial and error must occur first. The species-specific sections (Section 5.0) will highlight enhancement techniques that have been used successfully in the Gulf region.

3.7.1 Polishing

Polishing involves using various grades of abrasive papers and polishing compounds to smooth the cut surface of the section. Large otoliths, embedded or not, can be polished with 400-800 grit wet-dry sand paper while larval and juvenile embedded otolith sections are typically polished with 1000-1500 grit. Electric polishers, gem polishers, buffing wheels, and hand polishing have all been used to remove saw marks and other surface imperfections. Alternatives to polishing include: covering or coating the surface with clove, cedar, or immersion oil, glycerin, Flo-Texx, or Loctite (Figure 3.49). These solutions reduce light refraction making ring identification easier to




Figure 3.49. First dorsal spine from a tripletail, *Lobotes surinamensis*, viewed in clove oil.

the reader (Section 3.7.4). **Note: Prolonged exposure to clove oil, cedar oil, or glycerin will result in reduced readability and should be used with caution (see Section 3.7.4).**

3.7.2 Etching

Acid etching is a technique commonly used to enhance otolith microstructure, especially daily growth rings. This technique is also employed when otoliths contain growth zones or rings that are either too small or too faint to obtain accurate counts. This method takes advantage of the differing chemical composition of the opaque and translucent zones of the otolith by application of a chemical that will differentially dissolve the organic and inorganic components within the matrix (Pannella 1980). The chemical is most often an acid solution applied to an otolith thin section that will dissolve the regions of concentrated organic material (the translucent zone), more so than the calcified opaque zone. Three solutions used for etching by Davis et al. (1988) include immersion in 0.1 M disodium salt EDTA for 15 to 20 minutes, immersion in 1% HCl solution for 20 to 30 seconds, or immersion in 2% Histolab RDO for five minutes. The etched sections are then viewed under a Scanning

Electron Microscope (SEM) (Figures 3.50A and B). An alternative to viewing the section is to create a replica of the etched surface using an acetate peel. The majority of otolith studies in the Gulf states focus on species that do not require the use of etching for analysis.




Figure 3.50. A.) Cross section of *Gymnothorax sp.* (moray eel) leptocephalus (larval) otolith, to show growth rings, SEM, X 4,930. B.) SEM micrograph of medial portion of a whole swordfish otolith, *Xiphias gladius*.

3.7.3 Staining

Similar to the application of oils or clearing substances, stains may be used to enhance the contrast between opaque and translucent growth zones, and more clearly define external and internal microstructure of the otolith. Dyes for this purpose generally act in one of two ways: 1) differential diffusion (uneven staining) of the protein and calcium matrixes or 2) reaction solely with the calcium carbonate portions of the otolith (Gauldie et al. 1998). Histological stains are most effective, and commonly used stains include Alizarin Red, Aniline Blue, Crystal Violet, and Toluidine Blue; the darker colors prove to be more effective (Richter and McDermott 1990). It is recommended that otoliths (whole sections) be exposed to the dyes from a minimum of one hour to as long as several days. Previous research by Richter and McDermott (1990) demonstrates that success in staining requires trial and error with different stains based on the properties inherent to the otolith of the individual species. Variance in the effectiveness of dyes between samples is likely due to interspecific differentiation in the otolith's proteinaceous otolin composition impacting the absorption of the stain and its reactivity with the section's surface. Staining works best when combined with other enhancing techniques such as acid etching (acidification of the stain), thin sectioning, and use of transmitted light, and has been demonstrated as an effective enhancement procedure (Gauldie et al. 1998, Richter and McDermott 1990, Albrechtsen 1968, Bouain and Siau 1988). Staining is often successful when used to aid in interpretation of otoliths that exhibit indistinct growth zones or annuli such as Florida pompano, *Trachinotus carolinus* (K. Guindon, FWRI, personal communication).

3.7.4 Clearing

Clearing an otolith section refers to the

process of soaking a whole otolith or otolith section in a fluid medium that facilitates the passage of light through the specimen. It is used for: 1) a reduction in the appearance of saw marks and other surface imperfections with the application of an oil, glycerin, alcohol, or water (temporary) or 2) the perfusion of the clearing medium into growth zones within the section (permanent). Clearing, in this section, will refer to the perfusion of the clearing medium into otoliths microstructure by soaking the whole otolith in either clove oil, cedar oil, or glycerin. The duration of soaking is critical in achieving good contrast; however, once applied, the effect can continue and eventually render a section unreadable. Therefore, caution must be exercised when attempting this technique as time of soaking is dependent upon objective, species, and the otoliths size.

The soaking media effectively saturates the protein between the calcium carbonate crystals. Clearing usually affects the summer growth zone first. Continued soaking will eventually clear the opaque zones and eliminate any contrast optically washing out growth increments in the entire section/otolith. Therefore, careful removal of the clearing media must occur before long-term storage of a section.

3.7.5 Baking

Baking otoliths (whole and sections) is a technique adapted from the "Break and Burn" methodology (see Wischniowski and Bobko 2000 for a complete description). Although baking works very well for certain species, it may not with others, and considerable trial and error is involved initially. Variation of oven temperatures, baking times, and oven types will yield considerable differences in the outcome of the method. Baking time is generally a function of otolith size, desired final color, and is very subjective (Figure 3.51). The advantage



Figure 3.51. Baked otoliths in tray.

of baking over burning is that the outer margin is not scorched beyond a readable state. At this time, baking has been used with limited success on a few species in the Gulf.

3.7.6 Filters

Several filters are available through microscope vendors and scientific suppliers that can alter the light source being used to interpret marks on otolith, spine, or finrays. Polarization is commonly used throughout the Gulf states to enhance ring identification. Color filters have also been used with moderate success for particular species (Figure 3.52).

3.8 Microscopy, Image Analysis, and Measurements

Otolith sections can be viewed under a low-power or stereomicroscope using reflected light, transmitted light, or a combination of the two. The choice of reflected or transmitted light is often made based on the preference of the reader, but subtle differences in readability may occur between illumination types (see Section 4.0 for discussion).




Figure 3.52. Cross-section of the dorsal fin-rays from an age-6 white grunt (*Haemulon plumieri*) viewed with a green filter (540nm narrow-band).

In recent years, the magnified image of otolith sections have been digitized, viewed, and analyzed using image processing software packages that utilize frame-grabbers and analog or digital cameras. This allows the scientist to



Figure 3.53. Image analysis station used to interpret an otolith section. System includes microscope, camera, computer, monitor, and interpretation software.

acquire an image of otolith sections, view it on a video or computer monitor, recognize and mark the core and rings, and measure distances from the core to each ring, the core to the edge, and between rings (Figure 3.53). Ring counts and distance measurements are then immediately stored in a computer file eliminating transcription errors that can occur if image measurements are manually recorded and entered into a computer. Most of these software packages allow the reader to enhance the saved image making annuli recognition easier. Some of the more advanced packages can even automate the otolith reading process by guiding the reader through the entire process. Image analysis is also beneficial in that two or more scientists can discuss the features of otolith sections without having to look into a microscope. This allows for quick resolve of differences between readers.

4.0 Age Determination

This section is designed to give the reader guidance in age interpretations using otoliths. We have used a common sciaenid otolith as the model because its features are typically clear and obvious. Other species' otoliths can be more difficult to interpret and several species-specific accounts are listed in Section 5.0.

Throughout Section 4.0, an example data sheet is provided to track the procedure as the otolith is processed and an age determined for a fish with a July birthdate (Figure 4.1). This data sheet is purely for illustration but indicates the minimum data that should be recorded.

Fish Id.	Capture Date	# Rings	Margin Code	Biological Age	Age Group
ST00001	06/03/2001				
ST00002	06/03/2001				
ST00003	07/14/2001				



Figure 4.1. Example datasheet and section prior to assignment of rings, margin code, or age.

The appearance of structures used to age fish will vary under different illumination methods. Transmitted light (light from below passed upward through the section) and reflected light (light from above) will produce opposite contrasts in the observed ring patterns and the terminology used to describe the images

can often be confused if the light source is not specified. That is why it is important to record the light source used when interpreting structures. Transmitted light (Figure 4.2A) makes the image appear as alternating wide (light) and narrow (dark or amber) rings while reflected light (Figure 4.2B) reverses the appearance. Either illumination method is useful and merely a personal preference. **However, for consistency in this section, the use of transmitted light is assumed unless stated.**




Figure 4.2. Otolith section viewed under A.) transmitted light where opaque zones appear dark and B.) reflected light where opaque zones appear light.

4.1 Otolith Development

A basic understanding of otolith

development through successive periods of otolith ring formation is necessary to interpret the information contained in the structure. An otolith contains annual growth zones, each made up of a translucent and an opaque “ring” or zone. In the southern U.S., the translucent ring is usually wider than the opaque ring and represents a period of faster growth (summer). The opaque ring is usually deposited during slower growth (winter) and is relatively narrow (see Section 2 for a detailed description on ring formation). This increment, which includes a single translucent and opaque ring, is an annual growth increment.

The exterior surface of a whole otolith may reveal observable rings. While some of these rings correspond with opaque rings observed in sectioned otoliths, it is **not** always the case (Section 3.6.4). The savings in time and effort of being able to enumerate rings on a whole otolith is obvious and tempting; however, validation is necessary to verify that rings counted on a whole otolith represent the number of rings that are observed in sections. For example, rings counted on whole striped mullet (*Mugil cephalus*) otoliths in Mississippi were consistently one ring fewer than the number counted on sections (Figure 4.3A and 4.3B).

While binocular dissecting microscopes

yield the clearest view, more advanced image analysis systems can be used. An analog or digital video camera attached to a microscope and a television or computer monitor allow multiple individuals to view the same image at one time. By attaching the video camera to a frame grabber card installed in a computer the images can be saved, annotated, and cataloged or archived. This system can be further enhanced by installing image analysis software that gives the user the ability to enhance the otolith images and perform various analytical and quantitative tasks, such as measuring inter-annular distances on the otolith. Image analysis systems have also been used to rapidly enumerate measurements used to back-calculate the length at ring development and automatically determine number of rings on the otolith.

4.2 Ring Enumeration

Counting opaque rings in an otolith may seem straightforward, but for some species, separate opaque rings are not distinct. Two specific problems can be encountered: 1) identifying the location of the first opaque ring near or within the core, and 2) an opaque ring beginning formation very near or on the edge of the otolith. If the timing of opaque ring deposition is concurrent with or immediately following spawning, the first opaque ring may




Figure 4.3. Rings observed in a mullet otolith using A) a thin section and B) a whole otolith.

be hidden within the core region. If time of capture is concurrent with ring deposition, a distinct ring may or may not be observed at the otolith's margin. When rings are not particularly clear, techniques can be used to help discern rings and are discussed separately within each species account when they apply (Section 5).

Ring enumeration and edge development are typically made along the sulcus from the center of the core to a selected position on each ring, such as the midpoint, and to the otolith margin (Figure 4.4). The number of opaque rings are counted and recorded next to the corresponding fish identification number. These ring counts should be 'blind readings' meaning without any knowledge of fish size or capture date. A second enumeration should be made by another, independent reader. This is commonly referred to as 'verification.' Consensus is achieved by revisiting enumeration disparities

Fish Id.	Capture Date	# Rings	Margin Code	Biological Age	Age Group
ST00001	06/03/2001	3			
ST00002	06/03/2001				
ST00003	07/14/2001				





Figure 4.4. Highlighted core and subsequent opaque rings on an otolith section with the sulcus designated in red.

between readers or by a third party. A final ring count is then recorded for each fish.

4.2.1 Margin Codes

Another necessary step when assigning ages to fish entails describing the relative stage of ring formation on the outer edge of an otolith's margin. Code 1 is assigned to the presence of an opaque ring at the edge and codes 2, 3, and 4 are assigned to progressive development of the translucent ring at the edge (Figure 4.5). Using the monthly frequency of occurrence of Code 1 through a calendar year can validate whether the formation of the opaque ring occurs on an annual basis (Figure 4.6). The determination of which 'third' the translucent ring has completed is somewhat subjective; however, the presence/absence of the opaque ring is relatively straightforward. **The relative interval distance between**



Code 1.	opaque zone present on edge
Code 2.	translucent zone forming to 1/3 complete on edge
Code 3.	translucent zone 1/3 to 2/3 complete on edge
Code 4.	translucent zone 2/3 to fully complete on edge

Figure 4.5. Codes identifying proportional margin development on sectioned otolith.

rings changes as the fish ages, owing to the geometry of the otolith and the rate of growth represented in a given annual growth zone. Translucent and opaque rings usually become progressively narrower further from the core (Figure 4.7). The distances observed in the completed ring(s) closest to the edge are those used to judge the outer margin or proportion of completion of the outer ring being evaluated. Multiple codes can be observed in different fish captured at the same time because the timing and duration of ring development can be protracted over several months.




Figure 4.6. Frequency of occurrence of margin Code 1 over twelve months, or on an annual basis.

Timing of initial deposition of opaque material at the edge of an otolith and subsequent completion of the opaque ring for a particular year may take a relatively short period of time (one to two months) for an individual fish (Figure 4.8). When observing this same process over a large population, the time between the first evidence of deposition in some fish until all fish are exhibiting translucent deposition (opaque deposition has ceased), may be as long as five to six months. In addition, the actual timing of formation is not necessarily concurrent with a birth date. Once determined, the margin code must be recorded (Figure 4.9).




Figure 4.7. Change in relative distance, or narrowing of translucent area, for each progressive growth zone.

4.2.1.1 Quality of Otolith Section Margins

When viewing otolith sections, the reader may initially interpret an opaque region at the margin (appearing dark if using transmitted light and opaque white under reflected light) as a growth annulus if the section was not cut absolutely perpendicular to the core (Figure 4.10). In these cases, if the microscope slide is slowly tilted by hand or flipped over and viewed from the other side, the front and back planes of the section are realigned, creating a perpendicular margin relative to the viewer. The perceived opaque zone caused by an oblique view of the edge will disappear as the sharp edge of the section is presented. While off-angle sections are a very common and routine occurrence, not all new readers are aware of the phenomenon. These same technique can be used to correct annulus double images in the otolith section interior region.

4.3 Assignment of Age

The analysis has now provided a **ring count** and a **margin code**. Both of these parameters have been obtained by physically viewing the otolith, understanding/recognizing what the rings are, counting the rings, observing the margin or edge, and recording that data.

Fish Id.	Capture Date	# Rings	Margin Code	Biological Age	Age Group
ST00001	06/03/2001	3	2		
ST00002	06/03/2001				
ST00003	07/14/2001				




Figure 4.8. Mean margin increment distance plotted over a 20 month period indicating that opaque ring formation begins in February.




Figure 4.9. Break out section of otolith edge used for margin code assignment.

Biological age and age group are then assigned from these data, taking into account the timing of opaque ring formation, date of capture and an estimated hatch date or birthday. The following discussion gives generalized examples to illustrate the concepts that are applied to these data to arrive at a useful age for each fish.




Figure 4.10 An illustration of a mounted section from an age-5 fish. Because the otolith was sectioned slightly off-axis or off-plane, the section appears to have six annuli but actually only has five. The last ring is the bottom edge of the section visible through the thin section.

4.3.1 Biological Age

Because ring formation and birthdate may not coincide, the number of rings observed on an otolith is **not necessarily** the fish's age in whole years. In reality, the age of a fish in whole years and the number of rings coincide only during one month (time/period) per year. During all other months the age of the fish is the number of rings, plus or minus the time (months) before or after its closest birthday.

An example would be the fish with a July birthday that has just finished forming its third opaque ring in April and is captured June, but will not become three years of age for another month. All of this makes assigning an age to a fish more than just using the number of observed rings as the age of the fish. The method used to assign an age is dependent upon the ultimate use of the age data (Figure 4.11).

Fish Id.	Capture Date	# Rings	Margin Code	Biological Age	Age Group
ST00001	06/03/2001	3	2	2.9	
ST00002	06/03/2001				
ST00003	07/14/2001				



Figure 4.11. Example section with rings outlined ready to assign a biological age.

An age estimate and known length of the fish provides a basis for describing growth.

Having age determined with the greatest resolution would, in most cases, yield the most accurate and reliable estimates of growth. The ages assigned to fish for use in determining growth are called biological ages. Biological age could be defined as the time elapsed from birth to capture and can be expressed in months or converted to the nearest tenth of a year (for ease of mathematical manipulation; Figure 4.11).

An average hatch date can be estimated from fecundity data or from peak densities of larval/post larval fish (Figure 4.12A).




Figure 4.12. Birthdate determination using A. seasonal postlarval fish size and frequency data and B. seasonal Gonadal Somatic Index (GSI).

Estimates of mean time of spawning can be calculated by dividing the mean size of postlarvae at capture by an estimated daily growth rate; thus back dating to the time of spawning. Mean timing of spawning can be calculated from an indicator of spawning such as maximum gonadal somatic index (GSI) values (Figure 4.12B).

4.3.2 Assigning Age Groups

Stock assessments utilize cohort data as well as catch/population data grouped into ages. These data make up age groups representing single year classes or cohorts based on whole year ages. This grouping is needed to keep all fish sampled during a defined time period (calendar year, fishing year, etc.) together. While each year's offspring are considered a single cohort, there can be cohorts within the same year class as well. A good example of this is the bimodal spawning in spotted seatrout; two spawning peaks within one calendar year result in a spring cohort and late summer cohort. Therefore, we will use 'age group' rather than cohort to define the age (in whole years) of a fish at the time of capture. This age reflects the greatest age that the fish would have attained during the selected time period, typically a calendar year (Figure 4.13). This means that all fish which would attain age-1 would be assigned an age group-1, regardless of the biological age (month) when captured. This ensures that all fish within a cohort remain together when analyzing the age structure of a population.

An illustration of assigning number of rings, a biological age and age group to an age-0 fish, as it could be caught in any month over a calendar year, is shown in Figure 4.14. Number of rings are normally assigned at the time of reading. Biological age is assigned by evaluating the month of capture, number of rings observed relative to the month of opaque

ring formation and an estimated month of birth. The year group or cohort is assigned by determining the largest whole year age a fish will attain during a calendar (fishing) year.

Fish Id.	Capture Date	# Rings	Margin Code	Biological Age	Age Group
ST00001	06/03/2001	3	2	2.9	3
ST00002	06/03/2001				
ST00003	07/14/2001				



Figure 4.13. Example otolith section with all variables determined and biological age and age group assigned.

The impact of using these two different 'ages' on assembling an age structure is further illustrated in Figure 4.14. The age structures indicate a shift of younger fish into older age groups when using the year group method.

4.4. Quality Control in Processing

In production ageing of otoliths, several tests need to be conducted periodically to determine reader accuracy and precision of interpretation within individuals and between multiple readers. Additional training for processors in quality control should increase the acceptance of the science by managers and industry.




Figure 4.14. Timeline illustrating birth, birthdate, and periods of annulus deposition in an age-0 fish. Table illustrates the change in number of rings, biological age, and age class over one calendar year with a July 1 birthday.

4.4.1 Validation

As a general rule when working with a new species, it should **not** be assumed that opaque rings are annuli. Annual deposition of opaque rings must be ‘validated’ by any one of several methods.

4.4.1.1 Chemical Marking

The most direct method involves exposing a fish to tetracycline, calcein, or some other chemical that incorporates a mark on the otolith through a physiological process. Through release and recapture of this marked fish over time, one had a direct method for validating whether one opaque ring is deposited on an annual basis. A problem with this approach is that the potential for recapture can be low in open marine systems, making this method less practical. As an alternative, a marked individual can be held in captivity for an extended length

of time for validation. However, the timing of opaque ring deposition of a fish held in captivity may not reflect natural conditions in the wild and should be interpreted with caution.

4.4.1.2 Marginal Increment Analysis

Annual deposition of the opaque ring is more commonly validated by marginal increment analysis. The examination of the otolith edge condition for multiple fish captured over a time continuum (typically monthly) reveals the timing of formation of the last opaque ring. If opaque rings are found at the edge of the otolith only during one time period per year, it is inferred that the process is a yearly event (see Campana 2001 for review). Many times these data are presented as the monthly mean distance from the proximal edge of the last visible opaque ring to the margin of the otolith. Lowest monthly values of margin increments

observed during a calendar year reveal the timing of opaque ring deposition and if the minimum value is observed only once per year, it is inferred that the process is an annual event (Figure 4.9).

4.4.2 Accuracy

In practice, the accuracy of an age determination method may be known, but the accuracy of a particular set of age estimates is seldom known (Beamish and McFarlane 1995). So age validation commonly refers to validation of the method used to determine age. Validation of absolute age is rarely done and has been primarily accomplished through age determinations of recaptured, tagged fish after a long interval of time or through the use of radiocarbon or radiochemical methods compared to growth increment estimates (Campana 2001).

Validation is critical for initial age and growth characterizations of a given species and validation of absolute age should be the preferred goal, but it is often exceedingly difficult and so two steps are recommended (Campana 2001). First, determine the time and age when the first increment forms. It is commonly overlooked because it can be problematic. Second, verify the increment periodicity across the entire age range of interest such that annulus formation/increment periodicity is determined for young immature individuals and old mature individuals (not necessarily every age class). See Campana (2001) for a recent review and critique of validation approaches.

Assuming for a given species that initial age and growth characterization is complete, validation of increment periodicity has been accomplished, and there is consensus on interpretation of ageing structures, ageing programs can move into the production

phase whereby large numbers of samples are aged at regular intervals. At this stage, quality control monitoring becomes a very important component, including exchanges of age samples and cross-checking between laboratories (Boehlert and Yoklavich 1984, Morison et al. 1998).

4.4.3 Precision

As validation deals with error in accuracy, a second source of error that becomes critical in production ageing is precision or reader variability. Precision error is commonly reduced (improved) by resolving interpretation differences among readers. Precision errors often result in 'smeared' age distributions that tend to obscure strong or weak year classes. This interferes with attempts to track age-structure changes and to estimate mortality rates across time using an age-structured model, or when trying to compare age distributions with environmental or recruitment indices (Beamish and McFarlane 1995).

Some fish are difficult to age and precision errors are always inherent at some level, but experience is of key importance. There are a few, well documented approaches to quality control. Primarily they involve second readings or the use of a reference collection of resolved-age samples (Campana 2001). An example is the case whereby a primary reader may read all the otoliths and then an experienced secondary reader or tester may read a random sample of 20% without knowledge of the ages assigned by the primary reader. Examinations of bias and reader error (precision) estimates should be recorded and updated annually (Kimura and Lyons 1991).

4.4.4 Reference Collection

The use of reference collections serves many of the same purposes as reader-tester

comparisons and has potential advantages. The dominant use of reference collections are to test precision among readers and to monitor consistency in age interpretations over time. A reference collection allows monitoring of long-term drift, an increase or decrease in counts over time based on subtle changes in a reader's interpretation of the ageing structure. This cannot be accomplished as well with a reader-tester approach using contemporary samples (Campana 2001). A reference collection is also useful for training purposes (Campana 2001). A subset of the reference collection can be imaged and annotated and used to illustrate ageing structures and characteristics during the training of new readers.

The reference collection must be a set of prepared ageing structures for which known or consensus-derived ages are recorded. The idea is to incorporate prepared otoliths (not necessarily textbook examples) that are representative of all age/size groups, regions and collection sources likely to be encountered by readers. Furthermore, building the collection using samples collected year-round is encouraged to show all stages of margin or edge development. If year-specific differences are suspected, consider including samples from several years. Dry storage of the otolith preparations is recommended for long-term archiving rather than storage in solutions such as glycerine (Campana 2001).

Although the size of the collection is arbitrary, Campana (2001) recommends about 500 age samples per stock. This number is large enough to prevent memorization and allows subsets to be exchanged among different groups of otolith readers. A particular subset (i.e., 100) may be thoroughly documented and used as a training set. Over time the collection should be augmented as new materials and processing procedures are updated.

Production ageing programs have shown that following initial orientation and training, periodic tests of precision and bias using the reference collection will enable several readers to age with consistency (Morison et al. 1998, Campana 2001). Consistency among readers and over time is important even if the consensus-derived ages, which serve as a basis for age interpretation, are later found to be inaccurate. If this happens, re-interpretation of the reference collection would allow age corrections to be readily made to the historical data sets (e.g., see Stanley 1986).

A 'before and after' exercise is recommended for each ageing session and is important for both experienced and novice readers. In the case of an experienced reader, perhaps some time has passed since a given species was last aged (at least a year or two) and a subset of the reference collection needs to be re-aged to tune the reader and prevent drift. For the novice reader, a training subset should be aged until a sufficient level of precision is achieved and reader bias is minimized (Morison et al. 1998). Near the end of the ageing session, a reader-tester exercise should be conducted, where another sub-sample of the reference collection should be read blind (without knowledge of previous readings, dates, or fish sizes), in order to generate an estimate of precision for the session (see below).

4.4.5 Reader Comparisons

When readers compare age estimates in order to achieve consistency, they need to examine any biases such that one reader may tend to under- or over-age another. A good approach for graphically detecting bias is to plot pair-wise age comparisons or age-bias graphs (Campana et al. 1995a). For annual age comparisons, most workers estimate precision measures using either Average Percent Error

(APE, Beamish and Fournier 1981) or percent Coefficient of Variation (CV, Chang 1982). Both approaches are valid and one may be preferred for various reasons. Regression analysis has shown that either measure can be easily predicted from the other (Campana 2001). Care should be exercised that comparisons are made for similar values; either raw increment counts or final assigned ages. Because it may be common for readers to have subtle differences in edge interpretations that are often hard to resolve and can affect the increment count, final assigned ages would tend to yield lower precision errors. Increasingly, these measures of reader error (precision) are being incorporated directly into stock assessment models in order to statistically correct age-structure estimates (Richards et al. 1992, Beamish and McFarlane 1995, Crone and Sampson 1998). In practice, a measure of reader error would be used to adjust or correct a single set of age determinations. This equates to what would have happened if several readers had come to consensus on each age in the set.

4.5 Other Parameters and Their Usefulness

Fish growth is usually derived from plotting length against age and/or fitting those data to an equation that can be used to estimate length for a given age. Many times only larger/older fish are available for examination (i.e., large specimens of fish from fishing tournaments or dockside sampling of commercial catch). Size and bag limits may hamper collections of fish representing the full size range of the populations when using fishery-dependent data. The growth rates of younger year classes of fish species that can grow quite old is of interest when smaller, younger specimens are rarely encountered. These estimates can be compared to observed lengths for each given age and provide insight into the overall growth and survival of fish in the population. In these cases, lengths at age can be estimated from a

technique referred to as ‘back calculation.’ If the relationship of otolith radius versus fish length is linear, then an estimate of fish length relative to a location (ring) on the otolith can be calculated.

The linear relationship of otolith radius and fish length is validated by regressing a series of otolith radiuses against the fish lengths for fish that cover as many ages/lengths as possible. Obviously, if no young fish are available, fish covering all ages may be non-existent. Assuming the relationship is linear, lengths are then estimated for each age by the following formula:

$$L_e = D_r/D_m * L_t$$

where L_e = estimated length,
 D_r = distance from core to chosen ring,
 D_m = radius of otolith,
 L_t = total length of fish at capture.

This formula gives an estimate of length for each chosen ring. If each ring represents an annulus (i.e., ring one represents age-1), estimates of length can be calculated for several ages on each otolith, given the number of rings present. This method is called the ‘direct proportion’ method. Further refinement of the above formula includes the Y-intercept from the regression of total length and otolith radius, such that:

$$L_e = D_r/D_m * L_t + Y\text{-intercept}$$

where L_e = estimated length,
 D_r = distance from core to chosen ring,
 D_m = radius of otolith,
 L_t = total length of fish at capture.

This technique is commonly called the ‘Fraser-Lee’ or ‘modified direct proportion’ method and is used when the regression of fish length and otolith radius does not pass through the origin. This method adjusts for any somatic


length gained prior to otolith growth. Other similar methods have been used mainly with the intent of partitioning the variance into age effects and length effects. DeVries and Frie (1996) provide details of the above methods.

5.0 Species-Specific Otolith Characteristics and Processing Details

As noted in Section 3.3, the sectioning techniques used for each species will be determined by the equipment (i.e., sectioning saws) already available in a laboratory. Three saw styles are currently used around the Gulf region: the low speed wafering saw, the high speed wafering saw, and the high speed thin sectioning saw. Three methods of section preparation are currently used in the Gulf states: embedding whole otoliths in an epoxy resin, mounting a whole otolith to a glass slide, and free-hand cutting of whole otoliths followed by mounting on a slide for sectioning. Differences in fish shape and body size and otolith size among species require species-specific modifications to otolith extraction, preparation, and analysis. The following species accounts summarize these differences and highlight techniques currently being used in the Gulf region.

At the end of each species account, a timeline is provided for that species. The timeline includes the periods of spawning and annulus deposition for the species throughout its range from the published literature so those periods may appear wider than your specific location. Also included at the bottom of each timeline, is the monthly ring count, age designations, and cohort or year grouping over the first 2-3 years of the fish's life. The year grouping is used to assign a fish to a cohort (i.e. year class). For example, age in years when subtracted from the year of collection should equal the birth year. An accounting must be made for those fish that have not yet completed an annulus in the year of collection (usually by spring or summer), or, in some cases, completes an annulus early (before January 1st). Thus in those cases, one ring may be added or subtracted from the count. Correctly assigning an annual age, is an important task.

5.1 Red Drum *Sciaenops ocellatus*



Highlights

- Otoliths are large and relatively easy to locate and extract.
- Multiple sectioning techniques successful.
- Rings easily discernable.
- First distinct opaque ring forms at approximately 1.5 years of age.
- Long-lived species – up to 40+ rings.

Otolith Description

Red drum have large, stout sagittae that are thick enough to be opaque (Figure 5.1). The sagitta is slightly elongate and ovoid with a rather straight and slightly crenate dorsal margin and a convex ventral margin (Chao 1978). The anterior and posterior portions are about the same height, forming a rectangular surface. There are often one or more knobby protrusions on the distal face.



Figure 5.1 Proximal and dorsal views of red drum sagittal otolith.

The ostium of the sulcus is large and pear-shaped, and its expanded part does not reach the anterior margin. The 'J' shaped cauda of the sulcus acusticus is sharply bent, and its dorsal edge extends further into the ostium than its ventral edge. The rostrum and anterostrum are not distinguishable from one another. The core of the otolith usually lies just interior to the surface that faces outward from the midline of the fish. In the antero-posterior axis, the core lies adjacent to the junction of the ostium and cauda regions of the sulcus acusticus. The location of the otolith in the neurocranium is illustrated in Figure 5.2.

Otolith Extraction

Red drum otoliths can withstand expected impacts from otolith extraction devices without breaking. The otic capsule of red drum is somewhat convex, making it easy to identify through the gill cavity near the posterior base of the skull above the gills. It is relatively easy to cut away the surface of the exposed otic capsule with a heavy knife. In larger fish, otolith removal is best done using a hacksaw cut made




Figure 5.2 Location of red drum sagittal otoliths.

from the dorsal surface of the head to the otic capsule. Red drum otoliths are relatively robust across all life stages; however, due to the still fragile nature of young otoliths, extraction should be executed with care in smaller fish. Several different techniques are effective; some may be easier than others on different sized fish.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.3).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.3 Extraction of red drum otoliths through the top of the neurocranium.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule (Figure 5.4).
2. Bend the head of the fish forward to reveal the sagittae.
3. Remove the sagittal otoliths.



Figure 5.4 Meatsaw technique for extraction of otoliths from red drum.

Bottom Method

This method causes minimal visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic

capsule (Figure 5.5).

3. Chisel away the otic capsule to expose the sagitta.
4. Remove the otolith.
5. Repeat for the other side.



Figure 5.5. Extraction of otoliths from red drum through the operculum.

Otolith Processing

Due to the robust nature of this species, multiple techniques are acceptable and usually reflect available equipment. Generally, red drum sections are processed at approximately 0.5 mm. The following techniques have been used successfully throughout the Gulf.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long

- side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

High Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.1.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust load (1,000 g) and speed (3,000 rpm). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.
4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Red drum otolith sections call for special attention in the process of identifying the first annulus. Because red drum spawn in the late fall, just before the time of opaque zone formation, a dark zone is often visible around the core. However, the first distinct opaque mark is deposited during their second winter when the fish is about 14-18 months of age (Figure 5.6).

discouraged, as researchers have determined they yield highly inaccurate age data (Rohr 1964; D. Tremain, FWC, personal communication).





Figure 5.8 Mean monthly marginal increment (± 1 SD) for red drum in Florida waters with (a) one and (b) two annuli on otolith sections (from Murphy and Taylor 1990).

5.2 Spotted Seatrout *Cynoscion nebulosus*



Highlights

- Otoliths are large and relatively easy to locate and extract.
- Multiple sectioning techniques successful.
- Rings easily discernable.
- Distance from the core to the first opaque ring is variable.
- First ring formation occurs at <1 year.
- Generally fewer than 13 rings.

Otolith Description

Spotted seatrout have relatively large, elliptical, narrow sagitta that are opaque at most sizes (Figure 5.9). The dorsal margin is smooth and convex, whereas the ventral margin is slightly concave and crenelate (Chao 1978). The posterior portion of the sagitta is wider laterally.



Figure 5.9 Proximal and dorsal views of spotted seatrout sagittal otolith.

The sulcus acousticus is elongate with the ostium ovoid and the cauda long and bent with a short distal end. The marginal groove is distinct, and the rostrum and anterostrum are not distinguishable from one another. The otolith core lies just interior of the midline of the distal surface of the otolith and beneath the juncture of the ostium and cauda of the sulcus acousticus. The location of the otolith in the neurocranium is illustrated in Figure 5.10.

Otolith Extraction

Spotted seatrout otoliths are strong enough to withstand expected impacts from otolith extraction devices without breaking. They are easy to identify through the gill cavity near the posterior base of the skull above the gills due to the strongly convex surface of the otic capsule which is easily cut away with a heavy knife.

Several different techniques are effective; some may be easier than others on different sized fish.



Figure 5.10 Location of spotted seatrout sagittal otoliths.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.11).



Figure 5.11 Removal of the top of the cranium in small spotted seatrout.

2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule.
2. Bend the head of the fish forward to reveal the sagittae (Figure 5.12).
3. Remove the sagittal otoliths.



Figure 5.12 Meatsaw technique for otolith removal in spotted seatrout.

Bottom Method

This method causes minimal visible damage to the fish (Figure 5.13).

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.
3. Chisel away the otic capsule to expose the sagitta.
4. Remove the otolith.
5. Repeat for the other side.

Otolith Processing

Due to the robust nature of this species, multiple techniques are acceptable and usually reflect available equipment. Generally, spotted seatrout sections are cut to approximately 0.5 mm. The following techniques have been used successfully throughout the Gulf.



Figure 5.13 Removal of spotted seatrout otolith through the gill cavity.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

High Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.1.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust load (1,000 g) and speed (3,000 rpm). Make successive 0.5 mm cuts to

- obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.
4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Spotted seatrout have a protracted spawning season which may extend from April to September, depending on annual variation in climate (Figure 5.15). Ageing is fairly straightforward even though the location of the first annuli can vary widely in its distance from the core (Figure 5.14). Due to the protracted spawning season there may be a corresponding variation in age (months) at first opaque zone formation, which may occur from October through May depending on geographic location, but peaks around March and April (Figure 5.16).





Figure 5.14 Sagittal otolith section from an age-3 spotted seatrout.

5.3 Black Drum *Pogonias cromis*



Highlights

- Otoliths large and relatively easy to locate and extract.
- Multiple sectioning techniques successful.
- Rings easily discernable.
- First distinct opaque ring forms at approximately 1 year of age.
- Long-lived species – up to 55+ rings.

Otolith Description

Black drum have a robust otolith that is semi-circular in juvenile fish and becomes somewhat rectangular in mature fish (Figure 5.17). The otolith is opaque with an oblong



Figure 5.17 Proximal and dorsal views of black drum sagittal otolith.

ostium and a crescent-shaped cauda. The rostrum and anterostrum are not distinguishable from one another. The otolith core lies just interior to the midline of the distal surface of the otolith. Black drum sagittae are opaque in older juvenile and adult fish. The location of the otolith in the neurocranium is illustrated in Figure 5.18.

Extraction

Black drum otoliths are strong enough to withstand expected impacts from otolith extraction devices without breaking. The ventral surface of the otic capsule of black drum is somewhat convex, making it easy to identify through the gill cavity near the posterior base of the skull above the gills. It is relatively easy to cut away the surface of the exposed otic capsule with a heavy knife. A heavy bladed knife can also be used to cut from the dorsal skull base at about a 30 degree angle to the back of the ocular socket to open the



Figure 5.18 Location of black drum sagittal otoliths.

cranial cavity and expose the sagittae. In larger fish, otolith removal is best done using a saw cut made from the dorsal surface of the head to the otic capsule. This method can also be performed on smaller fish, but care must be taken that the cut does not extend through the otic capsule for risk of damaging the otoliths. Several different techniques are effective; some may be easier than others on different sized fish.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.19).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.19 Extraction of red drum otoliths through the top of the neurocranium.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule.
2. Bend the head of the fish forward to reveal the sagittae (Figure 5.20).
3. Remove the sagittal otoliths.



Figure 5.20 Meatsaw technique for otolith removal in black drum.

Bottom Method

This method causes minimal visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic



Figure 5.21 Extraction of otoliths from black drum through the gill cavity.

capsule (Figure 5.21).

3. Chisel away the otic capsule to expose the sagitta.
4. Remove the otolith.
5. Repeat for the other side.

Otolith Processing

Due to the robust nature of this species, multiple techniques are acceptable. The technique chosen will likely reflect your current equipment. Generally, black drum sections are processed at approximately 0.5 mm. The following techniques have been used successfully throughout the Gulf.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down, with the long axis parallel to the long side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

High Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.1.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust load (1,000 g) and speed (3,000 rpm). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-hand whole otolith sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.
4. Place slide in guide arm and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Ageing of black drum is relatively easy since opaque zones are normally very distinct (Figure 5.22). Black drum spawn in the winter at approximately the time of opaque zone formation; therefore, the first distinct opaque mark is deposited when the fish is about one year old (Figure 5.23).




Figure 5.22 Sagittal otolith section of an age-30+ black drum. Arrows indicate the first eight annuli.

For regional stock assessment purposes, three minimal parameters are recorded: number of rings, presence or absence of opaque ring at margin, and month of capture. Based on these three parameters, cohort and biological ages can be determined. Murphy and Taylor (1989) validated the timing of annuli deposition using marginal increment analysis (Figure 5.24). The acceptable birthdate for this species is April 1 (Figure 5.23).

Other Ageing Methods




Figure 5.23 Birthdate assignment timeline for black drum. Age and year group based on biological birthdate (April 1), number of rings, and January 1 to December 31 year.





Figure 5.24 Marginal increment analysis for black drum (from Murphy and Taylor 1989).

Whole black drum otoliths have not been used successfully in the Gulf region, and the usefulness of break and burn techniques for black drum has not yet been determined. However, this species may be a good candidate for break and burn.

Scales have been demonstrated to be useful in the first few years only. After age-3 annuli in scales become less consistent and resorption can occur at the core (J. Moran, ASMFC, personal communication).

5.4 Striped Mullet *Mugil cephalus*



Highlights

- Otoliths are relatively easy to locate and extract.
- Otoliths are fragile; care must be taken in removal.
- Generally one removal technique practiced.
- Multiple sectioning techniques successful.
- Rings relatively faint but discernable.
- First distinct opaque ring forms at approximately one year of age.
- Generally <8 rings.

Otolith Description

Striped mullet have small, fragile sagittal otoliths, which may break during extraction. The ventral surface is moderately crenate (Figure 5.25). The distal side is concave with the visible core lying in the center of the otolith. The sulcus runs along the proximal dorsal half of the otolith.



Figure 5.25 Proximal and dorsal view of sagittal otolith from striped mullet.

The posterior margin is rounded. The location of the otolith in the neurocranium is illustrated in Figure (5.26).




Figure 5.26 Location of striped mullet sagittal otoliths.

Otolith Extraction

Extraction begins by cutting the isthmus of the gill arch with a pair of angled head diagonal wire cutters (Figure 5.27). Next, gills may be



Figure 5.27 Cutting the striped mullet isthmus with wire cutters.

pushed aside or removed while bending the head back (dorsally) (Figure 5.28) and exposing the otic capsule (Figure 5.29). Caution should be taken on smaller specimens (<200 mm), because this action may rupture the otic capsule and expose or expel the sagittal otoliths. Insert a pair of wire cutters or chisel on the posterior section of the otic capsule and pry off the surface (Figure 5.30). Otoliths are small and



Figure 5.28 Striped mullet cranium forced upward exposing the posterior end of the otic capsule.



Figure 5.29 Removal of gill arches further exposes the otic capsule.

may become chipped or broken if care is not taken. For example, a striped mullet with a 280 mm fork length has an otolith 9 mm in length and 3 mm at its maximum width. Otoliths are removed with a pair of forceps and then rinsed with water (Figure 5.31). Samples are then dried and placed in coin envelopes or plastic zipper bags with pertinent information recorded on the outside.



Figure 5.30 opening the otic capsule with wire cutters.

Otolith Processing

Although this species tends to have relatively thin and fragile otoliths, each of the sectioning techniques described in Section 3.0 can be used with care.




Figure 5.31 Removal of the sagittal otoliths.

Low Speed Wafering Saw Technique

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

High Speed Wafering Saw Technique

Embedded Whole Otoliths (Section 3.4.1.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust load (1,000 g) and speed (3,000

- rpm). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

Note: Only use the grinder on small/fragile otoliths.

1. Firmly grasping the posterior end of the otolith, grind material until adjacent to the core.
2. Mount otolith half with core on labeled slide.
3. Holding slide in hand, grind down remaining material to approximately 1mm.
4. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Annuli in striped mullet are recognizable along the sulcus (Figure 5.32). The otolith radius and annuli are measured from the core at the base of the sulcal groove along a medial line adjacent to the sulcal groove. Annulus formation was validated by Thompson et al. (1989) in Louisiana waters, and generally




Figure 5.32 Sagittal otolith section of age-5 striped mullet. Black arrows indicate annuli. Note that the large opaque core is not counted.

begins in January and is complete by April. Striped mullet in the Gulf of Mexico are spawned around November-December (Thompson et al. 1989) and subsequently deposit a large opaque region around the core

through February (Figure 5.33). This may be regarded as the first winter mark. The second winter mark or first true annulus is generally located further from the core, because it is deposited when the fish are approximately 12 - 14 months of age. Illuminated from below, the opaque rings in the section are relatively well defined.

Other Ageing Methods

Scales were originally used for mullet ageing from the 1950s and have been used through the 1970s. Ibanez-Aguirre and Gallardo-Cabello (1996) compared scales and otoliths for ageing purposes and reported that scales could be used for young ages, but otoliths provided better resolution for the older age classes.





Figure 5.33 Birthdate assignment timeline for striped mullet. Age and year group based on biological birthdate (November 1), number of rings, and January 1 to December 31 year.

5.5 Southern Flounder *Paralichthys lethostigma*



Highlights

- Otoliths small, fragile, and comparatively difficult to locate and extract.
- Otolith pairs asymmetrical to each other.
- Left otolith recommended for sectioning.
- Multiple sectioning techniques successful.
- Rings discernable.
- First distinct opaque ring forms at approximately one year of age.
- Differential growth in males and females.
- Maximum validated age of eight years.

Otolith Description

Southern flounder sagittal otoliths have a flat arrowhead shape. As in numerous flatfish, southern flounder display morphological differences between right and left saggitae (Figure 5.34). The core of the left otolith is

located more posterior to center. Therefore, consistent use of the right or left otolith is recommended for ageing. The location of the otolith in the neurocranium is illustrated in Figure (5.35).



Figure 5.34 Proximal and dorsal views of southern flounder right sagittal otolith.




Figure 5.35 Location of southern flounder sagittal otoliths.

Extraction

Sagittal otoliths can be removed from southern flounder in two ways.

Top Method

This method requires the removal of the top of the neurocranium. The technique is the same as that used for other species even with the flounder's unusual anatomy (Figure 5.36).

1. Make a horizontal cut (parallel to the lateral line) just above the eye, back to the preopercle.
2. A vertical (dorsal) cut is then made intersecting with the first cut removing a triangular section of the fish's head, exposing the otic capsule and the otoliths within.
3. Right and left otoliths are easily removed with forceps.



Figure 5.36 Pop-the-top method for otolith removal in southern flounder.

Bottom Method

This method requires going through the gill cavity and is preferred when sampling a commercial catch as it minimizes visible damage to the fish.

1. Pull open the left gill cavity exposing the gills.

2. Using a chisel, scrape the gills back to expose the otic capsule (Figure 5.37).
3. Chisel away the otic capsule to expose the otolith.
4. Remove the left otolith with a forceps.
5. Repeat steps on right side.




Figure 5.37 Otolith removal from a southern flounder through the operculum.

Otolith Processing

Due to the small size of southern flounder otoliths, the technique of sectioning whole embedded otoliths appears to provide the highest quality sections. Because of the differences in the left and right sagitta, it is suggested that the left be used for sectioning and the right catalogued and stored for possible future use. Southern flounder otoliths should be cross-sectioned at a thickness of approximately 0.5 mm to obtain the best results.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the anterior/posterior axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.

4. Mount the core section and label appropriately.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

High Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.1.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust load (1,000 g) and speed (3,000 rpm). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Sectioning Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

Note: Only use the grinder on small/fragile otoliths.

1. Firmly grasping the posterior end of the otolith, grind material until core is visible.
2. Mount otolith half with core on labeled slide.
3. Holding slide in hand, section off remaining material.
4. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Opaque increments are easily distinguishable on both the dorsal and ventral sides of the sulcus in southern flounder otolith

cross-sections (Figure 5.38) as spawning and annulus deposition overlap for the most part (Figure 5.40). Ages are assigned based on opaque increment count and edge condition recorded as opaque or translucent using the criteria of Beckman et al. (1991) and on a birth date of January 1 (Wenner et al. 1990).



Figure 5.38 Sagittal otolith section from an age-4 southern flounder. Black arrows indicate annuli.

Validation of annual increments was reported using marginal increment analysis most recently by Fischer and Thompson (2004). Annulus deposition begins in the northern Gulf in January and is completed by the end of May (Figure 5.39).




Figure 39. Percentages of opaque margin edges are plotted against month of capture (Fischer and Thompson 2004).

Other Ageing Methods

Whole otoliths – Fitzhugh (personal communication) indicates that young southern flounder (age-0 to age-4) may yield good ages when read whole, but cautioned that

corroboration with sectioned otoliths must be completed. MacNair et al. (2001) and Sipe and Chittenden (2001) both concluded that whole otolith ageing was adequate for young fish (to age-14 in California halibut, *Paralichthys californicus*, and age-4 in summer flounder, *Paralichthys dentatus*). Both of these studies compared whole otolith ages to sectioned ages in these two species of paralichthids.

Flounder otoliths may be too fragile and thin to achieve acceptable results using the break and burn technique.

Flounder scales were unsatisfactory for age determination, due to a lack of consistent markings (Palko 1984).





Figure 5.40 Birthdate assignment timeline for southern flounder. Age and year group based on biological birthdate (January 1), number of rings, and January 1 to December 31 year.

5.6 Gray Triggerfish *Balistes capriscus*



Highlights

- Otoliths very small and nearly impossible to locate.
- First dorsal spine commonly used for ageing.
- Spine stored frozen due to vascularization.
- False annuli can occur.
- Embedding not required.
- Focus deterioration in older fish can result in loss of early annuli.

Otolith Description

The otoliths of the gray triggerfish change their direction of accretion over time and do not contain annual marks (Figure 5.41) (Ofori-Danson 1989, Johnson and Saloman 1984, Escorriola 1991, Wilson et al. 1995, Hood and Johnson 1997).




Figure 5.41 Proximal view of gray triggerfish sagittal otolith.

In addition, the relative small size of the otoliths make them nearly impossible to extract. Therefore, estimates of age and growth in gray triggerfish have been reported by numerous scientists using annuli evident in the first dorsal spine rather than using otoliths. The location of the otolith in the neurocranium is illustrated in Figure 5.42.




Figure 5.42 Relative location of the sagittal otoliths in a gray triggerfish.

Spine Extraction

Removal of dorsal spines from gray triggerfish is relatively straightforward and can be applied to many species. See Section 3.6.4 for a detailed description of the following methodologies. **Note: Due to the fact that spines are vascularized, failure to freeze spines will result in rapid deterioration!**

1. Cut the membrane between the first and second dorsal spine toward the joint (Figure 5.43, line A).
2. After the membrane is cut, insert the knife into the condyle socket behind the first dorsal spine, and remove any connective tissue holding the spine in place.
3. Applying pressure to the spine, pull it forward until it 'pops' out of the socket (Figure 5.43, line B).




Figure 5.43 A) Cutting plane and B) direction of pull for removal of the first dorsal spine in gray triggerfish.

4. Cut any remaining skin separating the spine from the fish (Figure 5.44).
5. Place the spine in a small, labeled envelope and store in a freezer **until ready to section.**

Spine Processing

As noted in Section 3.5.3, a modified combination of methods can be used to process the first dorsal spine of gray triggerfish. In order to ensure a definitive margin on the



Figure 5.44 Removal of dorsal spine from gray triggerfish.

posterior lobes, remove the skin from between and covering the lobes. This will enable the production of a section with a smooth, readable, and measurable margin. **Note: If the section is made below the lateral groove, the annuli will be truncated and impossible read all the way around the spine.** Two techniques have been used in the Gulf for this species on both the thin sectioning machine and low speed wafering saws, although any saw should suffice.

Thin Section Machine (Section 3.6.4.1.1)

1. Cut the dorsal spine above the condyle freehand (Figure 5.45).



Figure 5.45 Cutting gray triggerfish spine freehand on thin sectioning machine.

2. Adhere the distal portion of the spine to a slide on the cut edge.
3. Mount slide in chuck and cut remaining spine leaving a section adhered to slide.
4. Adjust thickness of section on the grinding wheel.

Low Speed Wafering Saw (Section 3.6.4.2.2)

1. Adhere spine to slide attaching only the ends with thermoplastic (Figure 3.40).
2. Place slide in chuck and make successive 0.5 mm cuts.
3. Adhere sections to slide.

Age Determination

The summer and winter growth zones in a gray triggerfish spine section are translucent and opaque, respectively, opposite the pattern found in an otolith. These annuli radiate outward from the focus. The focus in a spine section is the main channel of vascularization for the spine. The spine radius is measured as the distance from the focus to the margin of one of the posterior lobes, as seen in Figure 5.46.

There are several occurrences of pseudoannuli or “false annuli” in gray




Figure 5.46 Generalized cross section of dorsal spine.

triggerfish spines (Figure 5.47). False annuli associated with checks and zones that are

somewhat incomplete and irregular are usually found only in one part of the structure and often not in all structures. Although they are sometimes prominent, they are not associated with the growth zone that forms during the principal annual cessation or reduction in growth that produces the annulus (Casselman 1983). This problem can be corrected with the validation of the hard part. Although the cause is not known, it is believed they may be related to both larval settlement (false annuli near the




Figure 5.47 Cross section of an age-7 gray triggerfish spine indicating the core, radius, and annuli. False annuli occur where two annuli appear with a single dash.

focus) and adult spawning events (midsummer) (Ingram 2001). In addition, the first dorsal spine core can undergo resorption and become more vascularized, obscuring and even eliminating the first few zones in older fish (Figure 5.48) resulting in an underestimation of age (Casselman 1983).

After enumeration of the true annuli, estimate the biological age of the gray triggerfish by adjusting for a June-July spawning date in the northern Gulf of Mexico (Wilson et al. 1995, Ingram 2001); adjusting for an annulus formation date between January and April



Figure 5.48 Deterioration of the core region in the first dorsal spine of an old gray triggerfish.

For regional stock assessment purposes, three minimal parameters are recorded: number of rings, presence or absence of opaque ring at the margin, and month of capture. Based on these three parameters, cohort and biological ages can be determined.

Alternative Techniques

Since otoliths are not used to age gray triggerfish, break and burn would not be a useful alternative.

Work underway by Murie and Fioramonti (NOAA Fisheries - Panama City Laboratory, personal communication) indicates a high degree of success using finrays rather than spines to age this species (Section 3.6.4.2).

(Wilson et al. 1995) and adjusting for the date of capture (Figure 5.49).

Scales have not been used in this species successfully, due to the strong insertion of the scales into the triggerfish's tough skin (G.W. Ingram personal communication).




Figure 5.49 Birthdate assignment timeline for gray triggerfish. Age and year group based on biological birthdate (July 1), number of rings, and January 1 to December 31 year.

5.7 Red Snapper *Lutjanus campechanus*



Highlights

- Otoliths are ovate, laterally compressed.
- Otoliths are relatively easy to locate and extract.
- First increment can appear diffuse and difficult to discern.
- Opaque increment enumeration becomes increasingly difficult in older fish.

Otolith Description

Red snapper otoliths (sagittae) are large, ovate, laterally compressed, and exhibit an indented sulcus on the proximal surface (Figure 5.50). The rostrum and anterostrum are distinguishable. The location of the sagittae in the neurocranium is illustrated in Figure 5.51.



Figure 5.50 Proximal and dorsal views of red snapper sagittal otolith.




Figure 5.51 Location of sagittal otoliths in red snapper.

Extraction

Red snapper otoliths may break during contact with certain extraction tools. The otic capsule in red snapper is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernible once the gills have been removed or scraped back. The capsule surface is fairly thin, can appear transparent, and is relatively easy to chisel away.

Bottom Method

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market, as it minimizes visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.
3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.52).
4. Remove the otolith.
5. Repeat for the other side.



Figure 5.52 Removal of red snapper otolith through the operculum.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a

point below and behind the eyesocket exposing the brain (Figure 5.53).

2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.53 Extraction of red snapper otoliths through the top of the neurocranium.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule.
2. Bend the head of the fish forward to reveal the sagittae (Figure 5.54).
3. Remove the sagittal otoliths.



Figure 5.54 Meatsaw technique for extraction of otoliths from red snapper.

Processing

Due to the relatively large size of red snapper otoliths, multiple processing techniques are acceptable. The technique chosen will

likely reflect available equipment. Generally red snapper sections are processed at approximately 0.5 mm. The following techniques have been used throughout the Gulf.

4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Low Speed Wafering Saw Technique

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis (anterior-posterior axis) parallel to the long axis of the mold.
2. Locate the core and position block in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections onto slides.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.

Age Determination

Enumeration of annuli in red snapper otolith sections can be challenging to inexperienced personnel. The problem encountered most often by readers is determining the position of the presumptive first opaque increment nearest the core (Figure 5.55). Wilson and Neiland (2001) validated the annuli deposition using marginal increment analysis from December through June. Due to a protracted spawning




Figure 5.55 Section from the sagittal otolith of an age-3 red snapper showing first annuli as a diffuse opaque zone (reflected light).




Figure 5.56 Birthdate assignment timeline for red snapper. Age and year group based on biological birthdate (July 1), number of rings, and January 1 to December 31 year.

season (early May through late September) (Figure 5.56), there is assumed to be considerable variation in the distance from the core to the first opaque increment, which can appear as a diffuse ‘smudge.’ The increment may appear adjacent to the core region if the fish was spawned in the fall (Figure 5.57A) or may appear as an annuli outside the core if a fish was spawned in early summer (Figure 5.57B). The longevity of the species also increases the difficulty in obtaining accurate age estimates of older fish. After age-10, red snapper somatic growth slows dramatically




Figure 5.57 Transverse sagittal otolith sections of A) fall spawned and B) summer spawned red snapper (arrows indicate position of 1st increment).

and is reflected by a decrease in the accretion rate in the otolith. The opaque rings will appear

much closer together with distance from the otolith core (Figure 5.58).




Figure 5.58 Transverse section of sagittal otolith from an age-52 red snapper.

Other Ageing Methods

Bomb radiocarbon is a recent technique used to validate otolith age, which utilizes the increase in oceanic ^{14}C resulting from atmospheric testing of nuclear bombs during the 1950s and 1960s. Otolith section ages were validated through accelerator mass spectrometry analysis of bomb-produced ^{14}C in red snapper otoliths hatched before, during, and after the nuclear testing periods (Baker and Wilson 2001).

Break and burn has not been attempted on this species in the Gulf. Whole otoliths have not been used with any success. Scales have been unsuccessful after the first few years of age.

5.8 King Mackerel *Scomberomorus cavalla*



Highlights

- Otoliths are elongate, laterally compressed.
- Otoliths relatively easy to locate and remove.
- First ring may resemble a diffuse 'smudge' in section.
- Whole otoliths can be successfully aged up to age-6.
- Rings in sectioned otoliths are usually distinct in older fish.

Otolith description

King mackerel sagittae are small, elongate, laterally compressed, and have an indented sulcus on the medial side (Figure 5.59). The rostrum and antirostrum are easily distinguishable and extremely fragile. The location of the otolith is illustrated in Figure 5.60.



Figure 5.59 Proximal and dorsal views of king mackerel sagittal otolith.

Extraction

Otolith removal in king mackerel is relatively easy; therefore, any of the techniques




Figure 5.60 Location of the sagittal otoliths in king mackerel.

illustrated in Section 3.1 can be used. Due to the fishes size, the meatsaw technique is recommended when the condition of the head is not important. The otic capsule in king mackerel is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernible once the gills have been removed or scraped back. The capsule surface is fairly thin, can appear transparent, and is relatively easy to chisel away.

Bottom Method

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market as it minimizes visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.
3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.61).
4. Remove the otolith.
5. Repeat for the other side.



Figure 5.61 Removal of king mackerel otolith through under the operculum.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.62).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule.



Figure 5.62 Extraction of king mackerel otoliths through the top of the neurocranium.

2. Bend the head of the fish forward to reveal the sagittae (Figure 5.63).
3. Remove the sagittal otoliths.




Figure 5.63 Meatsaw technique for extraction of otoliths from king mackerel.

Processing

Sectioning preparation typically consists of embedding the otoliths in bullet molds (Section 3.3.1.3). In the Gulf, the primary sectioning apparatus used is the low speed saw, although the thin sectioning machine has also been used successfully. It should be noted that the NMFS Panama City Laboratory strongly recommends the use of the low speed saw for small otoliths such as the mackerels and suggests a comparison of the results from both types of saw before making a long-term equipment choice. For very young fish the otoliths can be read whole (see age determination below).

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

Note: Only use the grinder on small/fragile otoliths.

1. Firmly grasping the posterior end of the otolith, grind material until adjacent to the core.
2. Mount otolith half with core on labeled slide.
3. Holding slide in hand, grind down remaining material to approximately 1 mm.
4. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Whole Otoliths (Section 3.6.1)

With few exceptions, small king mackerel up to age-4 are much easier to age using whole otoliths. A good rule of thumb is to use whole otoliths to age males <80 cm FL and females <90 cm FL. The following is a brief methodology for ageing king mackerel using whole otoliths.

1. Place otolith, distal or concave side up, in watch-glass with water.
2. Use a dark stage and reflected light (preferably a fiber optic light) to view otolith.
3. Annuli are read on the distal side of the posterior half of the otolith; those in the corner formed by the posterior and ventral edges are often the easiest to identify.
4. Readability can almost always be improved by rotating the watch-glass and adjusting the angle and intensity of the light. Try illuminating the otolith through the side of the watch-glass if you have a fiber optic light.
5. Changing magnification, especially lowering it, will also improve readability on some otoliths.
6. Examine both left and right otoliths if available, as they often vary in readability.

In most cases the distance from the core to the first annulus will be much larger than all subsequent increments, although the increment between the first and second annuli will sometimes be quite large as well (Figure 5.64). If a whole otolith from a small fish seems especially difficult to read, try sectioning it, as occasionally the section will be more readable than the whole otolith, even in younger fish.

Ageing Sections

Annuli in sectioned king mackerel otoliths



Figure 5.64 Whole otolith from an age-2 king mackerel.

are almost always most readable in the dorsal portion, especially along the sulcal groove. With transmitted light and a compound microscope, all annuli except the first appear as fairly narrow dark marks (Figure 5.65). The first annulus is almost always the most difficult to identify, as it is often just a broad, diffuse




Figure 5.65 Otolith from an age-8 year-old king mackerel sectioned on a low-speed saw.

dark band. This first annulus sometimes is more apparent on the ventral portion of the otolith, even if subsequent annuli are not, so it always pays to examine that area if it is not clear on the dorsal end. One other time when the ventral portion should be examined is when the sectioned fish is very young (i.e., two or

three) as sometimes the annuli will be clearer there than on the dorsal portion. A common phenomenon in king mackerel otolith sections is for annuli to appear as doublets or couplets, which can lead to significant overageing problems if one is not careful. Adjusting the focus often helps resolve this problem. Another characteristic of these sections is that after the second or third annulus, the growth increments are almost always quite uniform in size, with little or no decrease in size with increasing age. Because of this trait, ageing older fish is no more difficult than ageing younger ones and suggests that otolith growth and fish growth seem to become decoupled in king mackerel at a fairly young age. Two techniques which may improve readability are using a polarizing filter and flipping the slide over on the microscope stage (this can make a big difference). If a section is very difficult to read and the fish is close to the minimum size for sectioning, examine the remaining otolith whole if available. Measuring increment distances from the core is somewhat problematic because the axis of growth in the otolith changes after the first ring is formed. Age determination in king mackerel is further complicated by its protracted spawning period (Figure 5.66) – May through October in the northern Gulf (Finucane et al. 1986) with a peak in September (Grimes et al. 1990). Annulus deposition occurs from March to May (Beaumariage 1973, Johnson et al. 1983). The oldest king mackerel aged to date was 26 years old (DeVries and Grimes 1997).

Other Ageing Methods

Break and burn is not recommended for this species due to the small size of the sagittal otoliths. Currently spines and other hard parts have not been attempted for this species, and no information exists on the use of scales for ageing king mackerel.





Figure 5.66 Birthdate assignment timeline for king mackerel. Age and year group based on biological birthdate (Sept 1), number of rings, and January 1 to December 31 year.

5.9 Greater Amberjack *Seriola dumerili*



Highlights

- Otoliths small and fragile, easy to break during extraction.
- Typically require embedding to section.
- Rings not always discernable requiring manipulation to read.
- Average life-span reported at 8-12 years, but as old as 15.

Otolith Description

Thompson et al. (1999) described greater amberjack sagittae as follows:

“Greater amberjack sagittae are small, thin, fragile and elongate in the anterior direction and bluntly crenelate at the posterior end. The medial surface is convex and has a deep, prominent sulcus. The anterior portion of the sagitta is curved laterally and the posterior end is relatively flat. The rostrum is longer than the antirostrum, but the difference increases with fish size. Prominent grooves and ridges are present on the lateral side of the sagittae and are nearly absent on the medial side” (Figure 5.67A and B).

The location of the otolith in the neurocranium is illustrated in Figure 5.68.



Figure 5.67 Proximal and dorsal views of greater amberjack right sagittal otolith.

Extraction

Otolith removal in greater amberjack is not easy. The otoliths are small and fragile, making it easy to damage them during extraction; however, while any of the techniques illustrated in Section 3.1 can be used, a few tend to be easier than others. The otic capsule in




Figure 5.68 Location of the sagittal otoliths in greater amberjack.

greater amberjack is located directly behind and under the brain making it difficult to get into through the gill cavity, although it can be done. The recommended approach is to cut through the head using the meatsaw technique or through the top of the neurocranium.

Bottom Method

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market as it minimizes visible damage to the fish, although it is difficult.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule (Figure 5.69).




Figure 5.69 Exposure of otic capsule in greater amberjack.

3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.70).
4. Remove the otolith.
5. Repeat for the other side.




Figure 5.70 Removal of greater amberjack otoliths after chiseling capsule open.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.71).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.71 Extraction of greater amberjack otoliths through the top of the neurocranium.

Larger Fish

1. Make a vertical cut in the skull at a point at the leading edge of the opercle between the brain and the otic capsule (Figure 5.72).
2. Carefully clean the cut to determine position relative to the otic capsule.
3. Some ‘digging’ may be required to locate the otic capsule; if necessary, another thin section can be cut to reach the capsules (Figure 5.73).
4. With great care, remove the sagittal otoliths.




Figure 5.72 Relative location of cut when sectioning greater amberjack head.



Figure 5.73 Otic capsules opened and sagittal otoliths exposed in posterior cross-section of greater amberjack head.

Processing

Sectioning preparation typically consists of embedding the otoliths in bullet molds (Section 3.3.1.2). In the Gulf, the primary saw which has been used is the low speed wafering saw, although the high speed wafering saw could also be used. The thin sectioning machine has been used successfully with this species using the freehand technique.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning
(Section 3.4.3)

Note: Only use the grinder on small/fragile otoliths.

1. Firmly grasping the posterior end of the otolith, grind material until adjacent to the core.
2. Mount otolith half with core on labeled slide.
3. Holding slide in hand, grind down remaining material to approximately 1 mm.
4. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

While difficult, greater amberjack can be aged when viewed in thin section. Annual deposition of opaque zones has been validated through marginal increment analysis (Manooch and Potts 1997, Harris et al. 2007) and using OTC-tagged fish that were subsequently recovered (Thompson et al. 1999). Spawning of greater amberjack occurs in the spring



Figure 5.74 Otolith section of an age-5 greater amberjack. White dots indicate annuli.

with both male and female GSI reaching a maximum in March and April (Murie and Parkyn 2008; Figure 5.75). Annulus deposition can occur from April to August, with the first annulus demarcated by a distinct translucent zone following the opaque core and core-ring or ‘smudge’ (Murie and Parkyn 2008; Figure 5.74). In these cases, the first readable annulus is actually deposited between 12 to 15 months (Figure 5.75).

Like many of the pelagics, the difficulty in ageing greater amberjack is due to the small size of the otolith. If the otolith is broken or damaged during extraction, age determination can be impossible. In addition, otoliths in this species, while not deformed, can lack any




Figure 5.75 Birthdate assignment timeline for greater amberjack. Age and year group based on biological birthdate (April 1), number of rings, and January 1 to December 31 year. A mark (ring or dark smear) can occur close to the core; however, the first true annulus does not occur until the fish is actually a year old.

evidence of rings at all; some otoliths just cannot be aged. While it is not practical to throw out difficult otoliths, it may be necessary at times for this species.

Other Ageing Techniques

Whole otoliths were not readable due to the lack of translucence even when immersed in clove oil or glycerin (Thompson et al. 1999). Break and burn is probably not practical due to the small size of the sagittal otoliths.

5.10 Spanish Mackerel *Scomberomorus maculatus*



Highlights

- Otoliths are elongate, laterally compressed.
- Otoliths relatively easy to locate and remove.
- First ring may resemble a diffuse 'smudge' in section.
- Whole otoliths can be successfully aged up to age-6.
- Rings in sectioned otoliths are usually distinct in older fish.
- Spanish mackerel generally do not live past age-11 on the Atlantic Coast.

Otolith Description

Spanish mackerel otoliths (sagittae) are small, elongate, laterally compressed, and have an indented sulcus on the medial side (Figure 5.76). The rostrum and antirostrum are easily distinguishable and extremely fragile due to their small size and the overall thinness of the entire otolith. The location of the otolith is illustrated in Figure 5.77.



Figure 5.76 Proximal and dorsal view of Spanish mackerel sagittal otolith.




Figure 5.77 Location of the sagittal otoliths in Spanish mackerel.

Extraction

Otolith removal in Spanish mackerel is relatively easy; therefore, any of the techniques illustrated in Section 3.1 can be used. Due to their small size, however, the meatsaw technique is not recommended. The otic capsule in Spanish mackerel is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernible once the gills have been removed or scraped back. The capsule surface is fairly thin, can appear transparent, and is relatively easy to chisel away.

Bottom Method

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market as it minimizes visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.
3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.78).
4. Remove the otolith.
5. Repeat for the other side.



Figure 5.78 Removal of Spanish mackerel otolith through under the operculum.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.79).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.79 Extraction of Spanish mackerel otoliths through the top of the neurocranium.

Processing

Sectioning preparation typically consists of embedding the otoliths in bullet molds (Section 3.3.1.2). In the Gulf, the primary saw which has been used is the low speed saw. For very young Spanish mackerel, otoliths can be read whole (see Age Determination below). The NMFS Panama City Laboratory strongly recommends the use of the low speed wafering saw when sectioning this species to ensure section clarity. It is suggested that a comparison of the results from both saws be made before making a long-term equipment choice.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.

3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

Note: Only use the grinder on small/fragile otoliths.

1. Firmly grasping the posterior end of the otolith, grind material until adjacent to the core.
2. Mount otolith half with core on labeled slide.
3. Holding slide in hand, grind down remaining material to approximately 1 mm.
4. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Whole Otoliths (Section 3.6.1)

With few exceptions, small Spanish mackerel up to age-3 are much easier to age using whole otoliths rather than sections. A good rule of thumb is to use whole otoliths to age males <45 cm FL and females <55 cm FL. It should be noted, however, that specimens as large as 60cm FL have been aged using whole and sectioned otoliths with high levels of agreement (J. Mareska, AMRD, personal communication). The following is a brief methodology for ageing Spanish mackerel using whole otoliths.

1. Place otolith, distal or concave side up, in watch-glass with water.
2. Use a dark stage and reflected light (preferably a fiber optic light) to view otolith.
3. Annuli are read on the distal side of the posterior half of the otolith; those in the

corner formed by the posterior and ventral edges are often the easiest to identify.

4. Readability may be improved by rotating the watch-glass and adjusting the angle and intensity of the light. Try illuminating the otolith through the side of the watch-glass if you have a fiber optic light.
5. Changing magnification, especially lowering it, will also improve readability on some otoliths.
6. Examine both left and right otoliths if available, as they often vary in readability.

In most cases the distance from the core to the first annulus will be much larger than all subsequent increments, although the increment between the first and second annuli will sometimes be quite large as well. If a whole otolith from a small fish seems especially difficult to read, try sectioning it. Occasionally the section will be more readable than the whole otolith, even in younger fish.

Ageing Sections

Annuli in sectioned Spanish mackerel otoliths are most readable in the dorsal portion, especially along the sulcus. With transmitted light and a compound microscope, all annuli except the first appear as fairly narrow dark marks. The first annulus is usually the most difficult to identify, as it is often just a broad, diffuse dark band (Figure 5.80). This first annulus sometimes is more apparent on the ventral portion of the otolith, even if subsequent annuli are not, so it always pays to examine that area if it is not clear on the dorsal end. One other time when the ventral portion should be examined is when the sectioned fish is very young (i.e., two or three) as the annuli will be clearer there than on the dorsal portion.

A common phenomenon in Spanish mackerel otolith sections is for annuli to appear as doublets or couplets, which can lead to significant overageing problems if one is

not careful. Adjusting the focus often helps resolve this problem. Another characteristic of these sections is that after the second or third




Figure 5.80 Sagittal otolith section from age-5 Spanish mackerel. White arrows indicate annuli.

annulus, the growth increments are usually uniform in size, with little or no decrease in size with increasing age. Because of this trait, ageing older fish is no more difficult than ageing younger ones and suggests that otolith growth and fish growth seem to become decoupled in Spanish mackerel at a fairly young age. Two techniques which may improve readability are using a polarizing filter on the light source and flipping the slide over on the

microscope stage (this can make a big difference). One other thing to try if the section is very difficult to read and the fish is close to the minimum size for sectioning is to examine the remaining otolith whole if available. Measuring increment distances from the core is somewhat problematic, because the axis of growth in the otolith changes after the first ring is formed.

Age determination in Spanish mackerel is further complicated by its protracted spawning period (Figure 5.81) – typically May through October in the northern Gulf (Powell 1975, Finucane and Collins 1986). Annulus deposition occurs during the spring or early summer (Powell 1975, Fable et al. 1987). The oldest Spanish mackerel aged by the NMFS Panama City Laboratory to date was age-11.

Alternative Techniques

Break and burn is probably not practical due to the small size of the sagittal otoliths, and the use of scales for this species has not yet been determined.





Figure 5.81 Birthdate assignment timeline for Spanish mackerel. Age and year group based on biological birthdate (Aug 1), number of rings, and January 1 to December 31 year. Early spawned fish can have a mark in the core region, but it is not generally counted as an annulus.

5.11 Atlantic Croaker *Micropogonias undulatus*



Highlights

- Otoliths relatively easy to locate and extract.
- Multiple sectioning techniques successful.
- Rings easily discernable.
- First distinct opaque ring forms at approximately 1.5 years of age.
- Generally less than ten rings.

Otolith Description

The sagittae in Atlantic croaker are very thick and shield shaped, often with a shelf or flange on the outer surface or on the dorsal margin (Figure 5.82). The ostium of the sulcus



Figure 5.82 Proximal and dorsal views of Atlantic croaker sagittal otolith.

is large, pear-shaped, and its expanded part does not reach the anterior margin. The 'J' shaped cauda of the sulcus acousticus is sharply bent, and its dorsal edge extends further into the ostium than its ventral edge. The rostrum and anterostrum are not distinguishable from one another. The core of the otolith usually lies just interior to the surface that faces outward from the midline of the fish. In the antero-posterior axis, the core lies adjacent to the junction of the ostium and cauda regions of the sulcus acousticus. The location of the otolith in the neurocranium is illustrated in Figure 5.83.

Otolith Extraction

Atlantic croaker otoliths can withstand expected impacts from otolith extraction devices without breaking. The otic capsule of Atlantic croaker is somewhat convex making it easy to identify through the gill



Figure 5.83 Location of Atlantic croaker sagittal otoliths.

cavity near the posterior base of the skull above the gills. It is relatively easy to cut away the surface of the exposed otic capsule with a heavy knife. In larger fish, otoliths can be removed using a hacksaw cut made from the dorsal surface of the head to the otic capsule. Atlantic croaker otoliths are relatively robust across all life stages, but due to the still fragile nature of young otoliths, extraction should be executed with care at smaller sizes.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eye socket exposing the brain (Figure 5.84).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.84 Extraction of Atlantic croaker otoliths through the top of the neurocranium.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule (Figure 5.85).
2. Bend the head of the fish forward to reveal the sagittae.
3. Remove the sagittal otoliths.



Figure 5.85 Meatsaw technique for extraction of otoliths from Atlantic croaker.

Bottom Method

This method causes minimal visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic

capsule (Figure 5.86).

3. Chisel away the otic capsule to expose the sagitta.
4. Remove the otolith.
5. Repeat for the other side.



Figure 5.86 Extraction of otoliths from Atlantic croaker through the operculum.

Otolith Processing

Due to the robust nature of the otoliths in this species, multiple techniques are acceptable and usually reflect available equipment. Generally, Atlantic croaker sections are processed at approximately 0.5 mm. The following techniques have been used successfully throughout the Gulf.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long

- side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

High Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.1.1)

1. Embed the whole otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust load (1,000 g) and speed (3,000 rpm). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.
4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Transverse otolith sections of Atlantic croaker show very clear, easily identified marks that can be used for aging. Typical sections have an opaque core surrounded by a blurred opaque band, composed of fine opaque and translucent zones (Figure 5.87). This band represents the first annulus. Because of Atlantic croaker's spawning season, the width of the first annulus varies among individual



Figure 5.87 Otolith section from age-8 Atlantic croaker. Arrows indicate annuli. Note: first annulus (approximately 3 months) is smudge near to core and not counted.

fish. Spawning typically occurs from August through November (Barbieri et al. 1994) with a peak in October (Holt et al. 1985), therefore the accepted birthdate for this species is October 1. Annuli deposition occurs from December through May (Figure 5.88) which was validated by Barger (1985) who reported that almost no

Atlantic croaker otoliths had annuli on the margins from June to November, but they all did by March (Figure 5.89). Late-spawned fish have a very narrow band that is almost continuous with the core; early-spawned fish have a wide, well-defined band clearly separated from the core. Because of this




Figure 5.89. Percent of otoliths with an annulus on the margin for Atlantic croaker otoliths (from Barger 1985).




Figure 5.88 Birthdate assignment timeline for Atlantic croaker. Age or year group based on biological birthdate (October 1), number of rings, and January 1 to December 31 year.

variation in width and proximity to the core, the first annulus is sometimes difficult to identify. Subsequent annuli are represented by easily identified, narrow, opaque bands that alternate with wider translucent bands outside the proximal margin of the first annulus.

For regional stock assessment purposes, three minimal parameters are recorded: number of rings, presence or absence of an opaque ring at the margin, and month of capture. Based on these three parameters, cohort and biological ages can be determined.

Other Ageing Methods

Whole otoliths have not been used successfully in the Gulf region. The usefulness of break and burn techniques for Atlantic croaker has not been determined; however, this species may be a good candidate for the technique. Atlantic croaker scales have not been demonstrated to be useful in the Gulf yet.

5.12 Sheephead *Archosargus probatocephalus*



Highlights

- Otoliths are ovate, laterally compressed.
- Otoliths relatively easy to locate and extract.
- Otoliths are relatively large, and multiple sectioning techniques can be used.
- Maximum age varies by region ranging from 14 yrs (FL), 20 yrs (LA), to 26 yrs (SC).

Otolith Description

Sheephead otoliths (sagittae) are relatively large, ovate, laterally compressed, and exhibit an indented sulcus on the proximal surface (Figure 5.90). The rostrum and anterostrum are easily distinguishable. The location of the sagittae in the neurocranium is illustrated in Figure 5.91.



Figure 5.90 Proximal and dorsal views of sheephead sagittal otolith.



Figure 5.91 Location of sagittal otoliths in sheephead.

Extraction

Sheepshead otoliths are not terribly fragile, but caution should be taken during extraction as they may break during contact with certain extraction devices. The otic capsule in sheepshead is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernible once the gills have been removed or scraped back. The capsule surface is fairly thin, can appear transparent, and is relatively easy to chisel away.

Bottom Method

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market as it minimizes visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.
3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.92).
4. Remove the otolith.
5. Repeat for the other side.



Figure 5.92 Removal of sheepshead otolith through under the operculum.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.93).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.93 Extraction of sheepshead otoliths through the top of the neurocranium.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule.
2. Bend the head of the fish forward to reveal the sagittae (Figure 5.94).
3. Remove the sagittal otoliths.



Figure 5.94 Meatsaw technique for extraction of otoliths from sheepshead.

Processing

Due to the relatively large size of sheephead otoliths, multiple processing techniques are acceptable. The technique chosen will likely reflect available equipment. Generally sheephead sections are processed at approximately 0.5 mm. The following techniques have been used throughout the Gulf.

Low Speed Wafering Saw Technique

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis (anterior-posterior axis) parallel to the long axis of the mold.
2. Locate the core and position block in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections onto slides.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long

- side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.
4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Enumeration of sheephead annuli in otolith sections is straightforward with the




Figure 5.95 Birthdate assignment timeline for sheephead. Age or year group based on biological birthdate (April 1), number of rings, and January 1 to December 31 year. A mark is occasionally formed near the core, but is not counted as an annuli.

exception of the first ring (Figure 5.95). The period of annulus formation in the northern Gulf is from March through May (Beckman et al. 1990), and spawning occurs offshore from February through April with a peak in March and April (Wilson et al 1988). The coincidence of ring formation and spawning can lead to dark cores in early spawners and opaque cores in late spawners (Figure 5.96A and B). In general, it is accepted that the core mark is not interpreted as a true annuli (Dutka-Gianelli and Murie 2001).

Other Ageing Methods


Break and burn has not been attempted on this species in the Gulf. Based on the size of the otolith, this technique may warrant further investigation. The ageing of whole sheephead otoliths has not been attempted in the Gulf. Scales have been used in the past to age

sheepshead, but when compared to otoliths, the use of scales was found to underestimate age by age-3.



Figure 5.96 Core region of A.) early spawned and B.) late spawned sheephead.

5.13 Gag *Mycteroperca microlepis*



Highlights

- Otoliths are large and relatively easy to locate.
- Multiple sectioning techniques have been used successfully.
- Annulus formation (opaque zone) is complete by spring to early summer.
- Rings easily discernable and identifiable even in whole otoliths up to age-8.
- Moderately long lived – up to 20+ rings although age-3 to 10 are most common.

Otolith Description

Like most of the groupers, gag otoliths (sagittae) are relatively large, laterally compressed and have an arrow shape (Figure 5.97). The rostrum, anterostrum, and sulcus are easy to distinguish and locate. It is not uncommon to see protrusions or irregularities along the ventral edge of the sagittal. The location of the otolith within the neurocranium is illustrated in Figure 5.98.



Figure 5.97. Proximal and dorsal view of gag sagittal otolith.




Figure 5.98. Location of the sagittal otoliths in the neurocranium of gag.

Extraction

Otoliths in large gag (> 14 years age) are heavy and robust. However, the otoliths are fairly thin and fragile in younger gag, so care should be taken during removal and storage. The otic capsule in gag is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernable, once the gills have been removed or scraped back.

Bottom Method

1. Cut the operculum to fold forward and open it wide out of the way.
2. Cut away the gill arches at their insertion.
3. Use a chisel to scrape away tissue from the otolith capsule. In grouper, and particularly in large specimens, the capsule tends to be beneath a hard and thick 'rise' of bony tissue and should easily be detected as a large knob or protrusion.
4. Pop open the capsule with the chisel and the relatively large sagittal otolith is easy to remove with forceps. (Figure 5.99)




Figure 5.99. Otic capsule exposed for removal of sagittal otolith through the gill cavity of gag grouper.

Top Methods

Meatsaw Technique

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule.
2. Bend the head of the fish forward to reveal the sagittae (Figure 5.100)
3. Remove the sagittal otoliths (Figure 5.101).



Figure 5.100 Vertical cut through the gag neurocranium in for sagittal otolith removal.



Figure 5.101 Exposed sagittal otolith being removed from vertical section of gag grouper head.

Processing

Due to the relatively large size of gag otoliths, multiple processing techniques are acceptable. As in other species, low-speed sectioning preparation typically consists of embedding the otoliths in molds. The use of a thin sectioning machine has also been very successful with this species and the approach is the same as for other species with large otoliths (e.g., red snapper, red drum). For younger ages, (approximately < age 8) annuli can often be readily counted from whole otoliths and age estimation is accomplished with better precision than from sections.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed. Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial transverse cut adjacent to the core.

2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled side.
4. Place slide in chuck and section off remaining material (to 0.5 mm).
5. Polish and apply mounting medium as preferred.

Age Determination and Validation

Whole Otoliths (Section 3.6.1)

1. Place otolith, concave side up (distal side up), in watch glass covered with water.
2. Use a dark stage (or dark watch glass) and reflected light source (preferably fiber optic).
3. Annuli are often most readily distinguished if the sagitta is tipped towards its dorsal edge using forceps (Figure 5.102).
4. By reading the annuli on the distal side, counting down along the tilted concave curvature to the dorsal edge at the deepest point (along the dorsal-ventral axis), good increment spacing is encountered. This area seems to facilitate making consistent counts and is the same area where transverse cuts would be made in older specimens. Adjusting the light source and the angle of tilt often helps to improve resolution of the annuli.
5. Upon making counts of approximately 8 annuli and greater, increment width and spacing appears to be very reduced and increments become difficult to resolve. At this point the otolith should be sectioned.




Figure 5.102 Dorsal edge and annuli on whole gag grouper otolith submerged in water.

In the whole otolith, the first and approximately 7 or 8 succeeding annuli are typically distinct. In sectioned otoliths, annuli are also relatively easy to distinguish and count in young fish but difficulty increases as older gag are encountered (i.e., >age 10). Multiple counting paths should be attempted in sections. But unlike several species in this manual, the best counting path for sections is often not along the sulcus, but rather along the dorsal or ventral margins (the same plane recommended for whole otolith annuli counts). This difference may be due to the relatively thin and laterally compressed nature of gag otoliths compared to many other species. Ageing methods and interpretations derived during earlier studies are still being used based on both sectioned and whole otoliths (McErlean 1963, Collins et al., 1987, Hood and Schlieder 1992, Johnson et al. 1993).

Whole and sectioned otoliths are assigned an age based on the count of annuli (opaque zones observed with reflected light) and the

degree of marginal edge completion. Typically, marine fish in the southeastern U.S. complete annulus formation (opaque zone formation) by late-spring to summer and this pattern seems to hold for gag (Figure 5.103). Annuli have been noted to form April to July in the Gulf (McErlean 1963, Hood and Schlieder 1992), but may take as long as August to complete in some areas or in some years (Collins et al. 1987). Spawning ranges from about December to May with peak spawning activity occurring February and March (Figure 5.103) (Hood and Schlieder 1992 and Collins et al. 1997).

Alternative Techniques

Spines and Rays

The method of spine and finray thin sectioning is actively being investigated as a non-lethal means of ageing gag. Ongoing work involving comparison of finray sections to sagittal otoliths is showing much promise.





Figure 5.103 Birthdate assignment timeline for gag grouper. Age and year group based on biological birthdate (March 1), number of rings, and January 1 to December 31 year. Although the core region has been observed with varying degrees of opacity, the first annulus occurs at around 0.8-1.3 years of age.

5.14 Red Grouper *Epinephalus morio*



Highlights

- Otoliths are ovate, laterally compressed.
- Otoliths are relatively easy to locate and extract.
- Whole otoliths can be successfully aged to 10 years.
- First annulus is typically clearly defined and identifiable.
- Maximum age 25-30 years with majority in 5-10 year age class.

Otolith description

Red grouper otoliths (sagittae) are thin, elliptical and concave (Figure 5.104). The posterior quadrant is greatly convoluted and irregular along that margin (Moe 1969). The relative location of the sagittae in the neurocranium is illustrated in Figure 5.105.



Figure 5.104. Proximal and dorsal view of red grouper sagittal otolith.




Figure 5.105 Location of sagittal otoliths in red grouper.

Extraction

Red grouper otoliths are somewhat thin and may break during contact with certain extraction tools, so care should be used during removal. The otic capsule in red grouper is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernible once the gills have been removed or scraped back. The capsule surface is fairly thin, can appear transparent, and is relatively easy to chisel away.

Bottom Method

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market as it minimizes visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.
3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.106).
4. Remove the otolith.
5. Repeat for the other side.




Figure 5.106 Removal of red grouper otolith through the operculum.

Top Methods

Smaller Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.107).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.



Figure 5.107 Extraction of red grouper otoliths through the top of the neurocranium.

Larger Fish

1. Make a vertical cut in the skull at a point just behind the centerline of the opercle through the otic capsule.
2. Bend the head of the fish forward to reveal the sagittae (Figure 5.108).
3. Remove the sagittal otoliths.

Processing

Considering the ease of reading both sectioned and whole red grouper otoliths, multiple processing techniques are acceptable. The technique chosen will likely reflect




Figure 5.108 Meatsaw technique for extraction of otoliths from red grouper.

available equipment. Generally red grouper sections are processed at approximately 0.5-0.7 mm. The following techniques have been used throughout the Gulf.

Low Speed Wafering Saw Techniques

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis parallel to the long axis of the mold.
2. Locate core and position block in chuck.
3. Adjust arm weight and speed.
4. Make successive 0.5 mm cuts to obtain the core region.
5. Mount the core sections.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.
4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Whole Otoliths (Section 3.6.1)

Whole otoliths are submerged in water, placed (distal or concave side up) in a black watch glass, and viewed through a stereomicroscope with the aid of reflected light from a fiber optic light source. Normally, whole otoliths are rolled back with the use of forceps to acquire a flat surface to age (Figure 5.109).

Identifying rings on whole and sectioned red grouper is straight forward. Moe (1969) and Johnson et al. (1998) documented that red grouper spawn from March through May, but may be as long as January to June. Deposition of the opaque band typically begins in



Figure 5.109 Whole red grouper read under low magnification submerged water of a black watch glass (Lombardi-Carlson et al 2008).

that otolith having a translucent edge on the margin between January 1 and June 30 should be advanced a year in age.

Alternative Techniques

Currently, no other techniques have been used in the Gulf to determine the age of red grouper. However, researchers elsewhere in the Caribbean have used urohyal bones (González et al. 1974, Valdés and Padrón 1980) and mesopterygoids (Rodríguez 1986). Rodríguez (1986) noted that two rings are deposited per year compared to otoliths in this species.

June and July (Figure 5.110). Lombardi-Carlson et al. (2008) determined





Figure 5.110 Birthdate assignment timeline for red grouper. Age and year group based on biological birthdate (May 15), number of rings, and January 1 to December 31 year.

5.15 Gray Snapper *Lutjanus griseus*



Highlights

- Otoliths smaller than other snappers, but similar in appearance.
- First annuli can be difficult to distinguish near core.
- Can live over 25 years but most of the fishery is under age-10.
- Can have check marks or false annuli associated in first year.
- Scales and otoliths have been used to age but scales are useful to age-12 at best.
- Whole otoliths have been used on smaller specimens.

Otolith Description

The gray snapper sagittae are much smaller than the red snapper, but are similarly ovate, laterally compressed, and exhibit an indented sulcus on the proximal surface (Figure 5.111). The rostrum and anterostrum are distinguishable, but quite fragile. The location of the sagittae in the neurocranium is illustrated in Figure 5.112.



Figure 5.111 Dorsal and medial views of gray snapper right sagittal otolith.

Extraction

Due to the small size of gray snapper otoliths, they may break during contact with certain extraction tools. The otic capsule in gray snapper is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernible once the gills have been removed or scraped back. The capsule surface is fairly thin, can appear transparent, and is relatively easy to chisel away. While gray snapper can reach 10 lbs and 24 inches, they are not generally large enough to warrant the ‘meatsaw’ approach for removal.

Bottom Methods

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market as it minimizes visible damage to the fish. Most of the gray snapper in the recreational and



Figure 5.112 Location of gray snapper sagittal otoliths in the neurocranium.

commercial catch can be sampled using this technique.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.



Figure 5.113 Removal of gray snapper otolith through the operculum.

3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.113).
4. Remove the otolith.
5. Repeat for the other side.

Top Methods

Smaller Fish

1. Make a cut from the front of the skull to a point just over and behind the eyesocket exposing the brain (Figure 5.114).
2. Remove brain to reveal the otic capsule and otoliths.
3. Remove the sagittal otoliths.



Figure 5.114 Lateral cut across top of gray snapper head exposing brain and otic capsules.

Larger Fish

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain.
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.

Processing

Although smaller than the other snappers, the gray snapper still has a relatively large otolith, which makes multiple processing techniques available. The best technique will likely reflect the equipment available in a lab. Generally gray snapper sections are processed at approximately 0.5 mm. The following techniques have been used in the Gulf.

Low Speed Wafering Saw Technique

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis (anterior-posterior axis) parallel to the long axis of the mold.
2. Locate the core and position block in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections onto slides.

Mounted Whole Otoliths (Section 3.4.2.2)

1. Mount whole otolith to slide, concave side down with the long axis parallel to the long side of the slide using thermoplastic.
2. Locate core and position slide in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

Gray snapper otoliths are more fragile than other snapper, therefore care should be taken when using the thin section machine, as a greater number of sections will chip or fracture.

1. Firmly grasping both ends of the otolith,

- make initial cut adjacent to the core.
2. Hand grind additional material until core is visible.
3. Mount otolith half with core on labeled slide.
4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination

Whole Otoliths (Section 3.6.1)

Whole otoliths are submerged in water, placed (distal or concave side up) in a black watch glass, and viewed through a stereomicroscope with the aid of reflected light from a fiber optic light source. Normally, whole otoliths are rolled back with the use of forceps to acquire a flat surface to age (Figure 5.115).



Figure 5.115 Whole gray snapper under low magnification in water.

Burton (2001) validated annuli deposition through age-9 on the east coast of Florida (Figure 5.116). Gray snapper spawn from April to November with a peak during the summer months in June and July, so it is assumed a June 1 birthdate (Figure 5.117).

The gray snapper is one of the smaller snappers, rarely exceeding 18 inches (45 cm) in length, and is almost always less than 10lbs. Maximum size is 24 inches and 10lbs. Sexual maturity is obtained after about 2 years of age, at lengths of 7-13 inches (18-33 cm) and the




Figure 2

Mean monthly marginal increments of gray snapper otoliths by age, areas pooled.

Figure 5.116 Gray snapper marginal increment analysis from Burton 2001.

estimated maximum age for this snapper is 25 years although Fischer et al (2005) recorded a 28 year old.

Other Ageing Techniques

Scales

Manooch and Matheson (1981) described the age of gray snapper using scales as well as whole and sectioned otoliths. Less than 20% of the scales were adequate for ageing. Whole otoliths were more useful than scales but the authors reported that the sectioned otoliths were "as legible as any we have seen."





Figure 5.117 Birthdate assignment timeline for gray snapper. Age and year group based on biological birthdate (June 1), number of rings, and January 1 to December 31 year.

5.16 Vermilion Snapper *Rhomboplites aurorubens*



Highlights

- As with other snappers, the first annulus near the core can be difficult to distinguish.
- Otolith is similar to red snapper but more fragile and must be sectioned thinner.
- Although the average age harvested is 4-5 years, they have been aged to 26 years.
- Size-at-age is highly variable in this species and not a reliable predictor of age.

Otolith Description

The sagittal otolith of vermilion snapper are relatively large, ovate, laterally compressed, and exhibit an indented sulcus on the proximal surface (Figure 5.118). The rostrum and anterostrum are distinguishable but quite fragile. The location of the sagittae in the neurocranium is illustrated in Figure 5.119.



Figure 5.118 Proximal and dorsal views of vermilion snapper right sagittal otolith.



Figure 5.119 Location of the sagittal otoliths in vermilion snapper.

Extraction

Vermilion snapper otoliths may break during contact with certain extraction tools. The otic capsule in red snapper is located near the posterior base of the skull behind the gills. The surface of the otic capsule is convex and easily discernible once the gills have been removed or scraped back. The capsule surface is fairly thin, can appear transparent, and is relatively easy to chisel away. Since vermilion snapper do not get very large, most of the methods will work. The meatsaw technique is acceptable but less necessary on this smaller species.

Bottom Method

The method of otolith extraction through the gill cavity is preferred when sampling a commercial catch intended for market as it minimizes visible damage to the fish.

1. Pull open the opercle to expose the gills.
2. Pull the gill arches back to expose the otic capsule.
3. Carefully chisel away the otic capsule to expose the sagitta (Figure 5.120).
4. Remove the otolith.
5. Repeat for the other side.



Figure 5.120 Removal of vermilion snapper otolith through the operculum.

Top Method

1. Make a cut from the back of the skull to a point below and behind the eyesocket exposing the brain (Figure 5.121).
2. Remove brain to reveal the otoliths.
3. Remove the sagittal otoliths.




Figure 5.121 Extraction of vermilion snapper otoliths through the top of the neurocranium.

Processing

Like most of the snapper, vermilion snapper have a relatively large otolith, which makes multiple processing techniques available. The best technique will likely reflect the equipment available in a lab. Generally vermilion snapper sections are processed at approximately 0.5 mm. The following techniques have been used in the Gulf.

Low Speed Wafering Saw Technique

Embedded Whole Otoliths (Section 3.4.2.1)

1. Embed the otolith with the long axis (anterior-posterior axis) parallel to the long axis of the mold.
2. Locate the core and position block in chuck.
3. Adjust arm weight (50-75 g) and speed (8-10). Make successive 0.5 mm cuts to obtain the core region.
4. Mount the core sections onto slides.




Figure 5.122 Whole vermilion snapper under low magnification in water.

Thin Section Machine

Free-Hand Whole Otolith Sectioning (Section 3.4.3)

Vermilion snapper otoliths are more fragile than other snapper, therefore care should be taken when using the thin section machine as a greater number of sections will chip or fracture.

1. Firmly grasping both ends of the otolith, make initial cut adjacent to the core. The grinding wheel can be used to grind down the post-rostrum to the core.
2. Hand sand additional material until core is visible, as the section can shatter if its ground to thin by machine.
3. Mount otolith half with core on labeled slide.
4. Place slide in chuck and section off remaining material.
5. Place slide into precision grinder arm and adjust caliper to 0.5 mm.

Age Determination and Validation

Whole Otolith (Section 3.6.1)

Whole otoliths are submerged in water, placed (distal or concave side up) in a black watch glass, and viewed through a stereomicroscope with the aid of reflected light from a fiber optic light source. Normally, whole otoliths are rolled back with the use of forceps to acquire a flat surface to age (Figure 5.122).

Annulus formation has been validated by several studies using marginal increment analysis (Beamish and McFarlane 1983, Zhao et al. 1997, Hood and Johnson 1999). Vermilion snapper spawning in the Gulf of Mexico occurs from May to September with some fish spawning several times a season (Nelson 1988, Hood and Johnson 1999). Zhao et al. (1997) noted that occasionally false annuli or checks were deposited close to the core that may have been the result of settlement or changes in feeding habitats. Those opaque zones in predictable distances were deemed as the true annuli.

Enumeration of annuli in vermilion snapper otolith sections can be challenging to inexperienced personnel. Due to a protracted spawning season (Figure 5.123), there is assumed to be considerable variation in the distance from the core to the first opaque increment, which can appear as a diffuse 'smudge.' The increment may appear adjacent to the core region if the fish was spawned in the fall or may appear as an annuli outside the core if a fish was spawned in early summer. The problem encountered most often by readers is determining the position of the presumptive first opaque increment nearest the core (Figure 5.124).




Figure 5.123 Birthdate assignment timeline for vermilion snapper. Age and year group based on biological birthdate (July 1), number of rings, and January 1 to December 31 year.

Additional Techniques

Break and Burn

At this time, no one has tried break and burn on vermilion snapper.

Scales

Scales and otoliths have been used to age vermilion snapper (Grimes 1978, Barber 1989). Scales are considered less reliable because as fish become older scales become more difficult to interpret (Grimes 1978, Collins and Pickney 1988) compared to otoliths and therefore discouraged.





Figure 5.124 Sagittal section (50X) from vermilion snapper with annuli (1-7) and opaque zones.

5.17 Gulf Menhaden *Brevoortia patronus*



Highlights

- Scales are used to age gulf menhaden.
- Scale rings are generally easy to discern.
- Sagittal otoliths are small and fragile; it is impractical to extract, process and read large numbers of whole or sectioned otoliths from gulf menhaden.
- Approximately 95% of gulf menhaden in the purse-seine catch for reduction are age-1 and age-2 fish combined.

Scale Description

The NOAA Fisheries Service laboratory at Beaufort, NC, has monitored the gulf menhaden purse-seine fishery for size and age composition of the catch since 1964 (Nicholson 1978). From the outset, program managers realized it was impractical to utilize otoliths to age gulf menhaden because; 1) sagittae were so small and fragile, and 2) large amounts of time and effort would be required to extract, process, and read whole or sectioned otoliths. Moreover, large numbers of ageing parts (around 10,000 or more) would be required to adequately characterize the fishery with annual landings of several hundred thousand metric tons. Thus, scales were selected as the ageing part of choice for gulf menhaden.

Chapoton (1967) determined that scale development on gulf menhaden began on larval specimens at around 21 mm FL and was complete in specimens > 27 mm FL. Gulf menhaden scales are generally thin and translucent.

Unlike most herrings, the posterior margin of gulf menhaden scales are pectinate (Figure 5.125). The anterior field is embedded in the integument. The entire scale is sculptured with fine circuli, which roughly parallel the anterior margin. The largest and most symmetrical (nearly rectangular) scales occur in a median




Figure 5.125 Scale from an age-2 gulf menhaden (188 mm FL, 142 g), showing the focus, scale edge, and first and second age rings.

lateral band above the lateral line and below the dorsal fin (Figure 5.126). Scale samples for ageing are removed from this area.




Figure 5.126 Area from which scale patch is removed for ageing gulf menhaden.

Scale Removal

A scale patch from gulf menhaden is removed with a blunt-edged scalpel. First, the scalpel is passed several times across the surface of the scales in the patch area (in an anterior to posterior direction) to remove excess slime and moisture. The scalpel is wiped clean, and inserted under the scales slightly posterior from where the patch will be taken (Figure 5.127). With a gradual move forward, several rows of scales are lifted from the body by pressing scales between the thumb and the scalpel (a scale patch of 20-30 scales).

The scale patch is placed in a small vial of water. A few drops of dishwashing detergent in the wash bottle, used to fill the vials, helps degrade residual slime on the scales.

Scale Processing

The scale patch is removed from the vial with recurved forceps and blotted dry on a paper towel. Scales are rubbed between the thumb and forefinger and/or middle finger to remove any residual integument. Individual scales are pulled from between the thumb and fingers, then mounted between two glass microscope slides. Ten scales (two rows of five) are placed on the first slide with pectinations pointing up, then they are covered



Figure 5.127 Scale patch ready to be removed with a scalpel from the flank of a gulf menhaden.

with the second slide. Slides are fastened together on the ends with short lengths of transparent tape. The cover slide is labeled with a unique port and specimen number combination (Figure 5.128).



Figure 5.128 Gulf menhaden scales from a single fish pressed between two microscope slides.

Age Determination

Nicholson and Schaaf (1978) found that ageing gulf menhaden with scales was problematic. They determined that most fish had well-defined scale rings, but others had no rings, or rings that were oddly spaced. Their criteria for scale ageing were based on appearance of the scales, number and spacing of the rings, and fish fork length at time of capture. Although admitting some subjectivity, they determined that fish with one or two scale rings displayed true annuli. For fish with oddly-spaced rings, it was possible to separate out age classes by ring location. Finally, for

Gulf menhaden spawn between October and April, with peak activity from December through March (Turner 1969, Fore and Baxter 1972). Scale annuli form in winter, and by convention the birth date for gulf menhaden is January 1 (Figure 5.130). Since the purse-seine fishery operates April through October, advancing ages because of calendar date (and unformed rings) is not an issue relative to the industrial fishery statistics.

6.0 Literature Cited

- Albrechtsen, K. 1968. A dyeing technique for otolith age reading. *Journal du Conseil Permanent International por l'Exploration de la Mer.* 32:278-280.
- Baker, M.S., Jr., and C.A. Wilson. 2001. Use of radiocarbon to validate otolith section ages of red snapper *Lutjanus campechanus* from the northern Gulf of Mexico. *Limnology and Oceanography.* 46(7):1819-1824.
- Barber, R. C. 1989. Age and growth of vermilion snapper (*Rhomboplites aurorubens*) in the northern Gulf of Mexico. M.S. Thesis. Texas A&M University. 87 pp.
- Barbieri, L.R., M.E. Chittenden Jr., S.K. Lowerre-Barbieri. 1994. Maturity, spawning, and ovarian cycle of Atlantic croaker *Micropogonias undulatus*, in the Chesapeake Bay and adjacent coastal waters. *Fishery Bulletin.* 92:671-685
- Barger, L.E. 1985. Age and Growth of Atlantic Croakers in the Northern Gulf of Mexico, Based on Otolith Sections. *Transactions of the American Fisheries Society.* 114:847-850.
- Beamish, R.J. and D.A. Fournier. 1981. A method for comparing the precision of a set of age determinations. *Canadian Journal of Fisheries and Aquatic Sciences.* 38:982-983.
- Beamish, R.J., and G.A. McFarlane. 1983. The forgotten requirement for age validation in fisheries biology. *Transactions of the American Fisheries Society.* 112:735-743.
- Beamish, R.J. and G.A. McFarlane. 1987. Current trends in age determination methodology. Pages 15-42, *In: R.C. Summerfelt and G.E. Hall (eds) Age and Growth of Fish.* Iowa State University Press. Ames, Iowa.
- Beamish, R.J. and G.A. McFarlane. 1995. A discussion of the importance of aging errors, and an application to walleye pollock: the world's largest fishery. Pages 545-565, *In: D.H. Secor, J.M. Dean and S.E. Campana (eds.) Recent Developments in Fish Otolith Research.* University of South Carolina Press.
- Beaumariage, D.S. 1973. Age, growth and reproduction of king mackerel *Scomberomorus cavalla*, in Florida. *Florida Marine Research Publication* 1. 45 p.
- Beckman, D.W., C.A. Wilson, and A.L. Stanley. 1989. Age and growth of red drum, *Sciaenops ocellatus*, from offshore waters of the northern Gulf of Mexico. *Fishery Bulletin.* 87:17-28.
- Beckman, D.W., A.L. Stanley, J.H. Render, and C.A. Wilson. 1991. Age and growth-rate estimation of sheepshead *Archosargus probatocephalus* in Louisiana waters using otoliths. *Fishery Bulletin.* 89:1-8.
- Blacker, R.W. 1974. Recent advances in otolith studies. Pages 67-90, *In: F.R. Harden Jones (eds) Chapter 4, Sea Fisheries Research.* John Wiley and Sons, New York.

- Boehlet, G.W. and M.M. Yoklavich. 1984. Variability in age estimates in *Sebastes* as a function of methodology, different readers, and different laboratories. *California Fish and Game*. 70(4):210-224.
- Bouain, A. and Y. Siau. 1988. A new technique for staining fish otoliths for age determination. *Journal of Fish Biology*. 32:977-978.
- Brothers, E.B., C.P. Mathews, R. Lasker. 1976. Daily growth increments in otoliths from larval and adult fishes. *Fishery Bulletin*. 74(1):1-8.
- Brothers, E.B. 1984. Otolith Studies. Pages 50-57, *In*: H.G. Moser, W.J. Richards, D.M. Cohen, M.P. Fahay, A.W. Kendall, Jr., and S.L. Richardson (eds.) *Ontogeny and Systematics of Fishes*. International Symposium (La Jolla), American Society of Ichthyology and Herpetology, Special Publication 1.
- Burton, M.L., 2001 Age, growth, and mortality of gray snapper, *Lutjanus griseus*, from the east coast of Florida. *Fishery Bulletin*. 99(2):254-265.
- Calliet, G.M. 1990. Elasmobranch age determination and verification: an updated review. NOAA Technical Report NMFS 90:157-165.
- Campana, S.E. 2001. Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *Journal of Fish Biology*. 59:197-242.
- Campana, S.E. and J.M. Casselman. 1993. Stock discrimination using otolith shape analysis. *Canadian Journal of Fisheries and Aquatic Sciences*. 50:1062-1083.
- Campana, S.E., and J.D. Neilson. 1985. Microstructure of fish otoliths. *Canadian Journal of Fisheries and Aquatic Sciences*. 42:1014-1032.
- Campana, S.E., M.C. Annand, and J.I. McMillan. 1995a. Graphical and statistical methods for determining the consistency of age determinations. *Transactions of the American Fisheries Society*. 124:131-138.
- Campana, S.E., J.A. Gagne, and J.W. McLaren. 1995b. Elemental fingerprinting of fish otoliths using ID-ICPMS. *Marine Ecology Progress Series* 122, 115-120.
- Campana, S.E., S.R. Thorrold, C.M. Jones, D. Gunther, M. Tubrett, H. Longerich, S. Jackson, N.M. Halden, J.M. Kalish, P. Piccoli, H. Depontual, H. Troadec, J. Panfili, D.H. Secor, K.P. Severin, S.H. Sie, R. Thresher, W.J. Teesdale, and J.L. Campbell. 1997. Comparison of accuracy, precision, and sensitivity in elemental assays of fish otoliths using the electron microprobe, proton-induced X-ray emission, and laser ablation inductively coupled plasma mass spectrometry. *Canadian Journal of Fisheries and Aquatic Sciences*. 54:2068-2079.

- C.A.R.E. (Committee of Age Reading Experts), Pacific Coast Groundfish Ageing Technicians. 1997. Manual on Generalized Age Determination Procedures for Groundfish. Under the sponsorship of Pacific States Marine Fisheries Commission. For: The Technical Subcommittee of the Canada/U.S. Groundfish Committee.
- Casselman, J.M. 1983. Age and growth assessment of fish from their calcified structures -- techniques and tools. *In*: E.D. Prince and L.M. Pulos (eds) Proceedings of the international workshop on age determination of oceanic pelagic fishes: tunas, billfishes, and sharks. NOAA Technical Report NMFS 8, 211p.
- Castonguay, M., P. Simard, and P. Gagnon. 1991. Usefulness of Fourier analysis of otolith shape for Atlantic mackerel (*Scomber scombrus*) stock discrimination. Canadian Journal of Fish and Aquatic Sciences. 48:296-302.
- Chang, W.Y.B. 1982. A statistical method for evaluating the reproducibility of age determination. Canadian Journal of Fisheries and Aquatic Sciences. 39:1208-1210.
- Chao, L.N. 1978. A basis for classifying western Atlantic Sciaenidae (Teleostei: Perciformes). NOAA Technical Report Circular 415.
- Chapoton, R.B. 1967. Scale development in the Gulf menhaden, *Brevoortia patronus*. Transactions of the American Fisheries Society. 96(1):60-62.
- Chilton, D.E., and R.J. Beamish. 1977. Age determination of lingcod (*Ophiodon elongatus*) using dorsal fin rays and scales. Journal of the Fisheries Research Board of Canada. 34:1305-1313.
- Chilton, D.E., and R.J. Beamish. 1982. Age determination methods for fishes studied by the Groundfish Program at the Pacific Biological Station. Special Publication of the Canadian Journal of Fisheries and Aquatic Science 60. 102 p.
- Christensen, J.M. 1964. Burning of otoliths, a technique for age determination of soles and other fish. Journal du Conseil Permanent International por l'Exploration de la Mer. 29:73-81.
- Collins, M.R., and J.L. Pinckney. 1988. Size and age at maturity for vermilion snapper (*Rhomboplites aurorubens*) (Lutjanidae) in the South Atlantic Bight. Northeast Gulf Science. 10(1): 51-53.
- Collins, M.R., C.W.Waltz, W.A. Roumillat, and D.L. Stubbs. 1987. Contribution to the life history and reproductive biology of gag, *Mycteroperca microlepis* (Serranidae), in the South Atlantic Bight. Fishery Bulletin. 85:648-653.
- Collins, L. A., A.G. Johnson, C.C. Koenig, and M.S. Baker Jr. 1997. Reproductive patterns, sex ratio, and fecundity in gag, *Mycteroperca microlepis* (Serranidae), a protogynous grouper from the northeastern Gulf of Mexico. Fishery Bulletin. 96:415-427.

- Collins, L. A., Fitzhugh, G. R., Lombardi, L. A., Lyon, H. M., Walling, W. T., and D. W. Oliver. 2002. Characterization of red grouper (Serranidae: *Epinephelus morio*) reproduction from the eastern Gulf of Mexico. Contribution Report 2002-07. Panama City Laboratory, Southeast Fisheries Science Center, National Marine Fisheries Service, Panama City, Florida. 10 p.
- Colura, R.L. and T.L. King. 1995. Using scale and otolith morphologies to separate spotted sea trout (*Cynoscion nebulosus*) collected from two areas within Galveston Bay. Pages 617-628, *In*: D.H. Secor, J.M. Dean, and S.E. Campana (eds.) Recent Developments in Fish Otolith Research. University of South Carolina Press, Columbia, South Carolina.
- Cowan, J.H., Jr., R.L. Shipp, H.K. Bailey, IV, and D.W. Haywick. 1995. Procedure for rapid processing of large otoliths. Transactions of the American Fisheries Society. 124:280-282.
- Crone, P.R. and D.B. Sampson. 1998. Evaluation of assumed error structure in stock assessment models that use sample estimates of age composition. Pages 355-370, *In*: Funk, Quinn, Heifetz, Ianelli, Powers, Schweigert, Sullivan and Zhang (eds.) Fishery Stock Assessment Models. University of Alaska Sea Grant College Program AK-SG-98-01.
- Davis, N.M., R.W. Gauldie, S.A. Crane, and R.K. Thompson. 1988. Otolith Ultrastructure of Smooth Oreo, *Pseudocyttus maculatus*, and Black Oreo, *Alloctytus sp.* Fishery Bulletin. 86(3):499-515.
- Dery, L.M. 1983. Use of laminated plastic to impress fish scales. Progressive Fish-Culturist. 45:88-89.
- DeVries, D.R., and R.V. Frie. 1996. Determination of age and growth. Pages 483-512, *In*: B.R. Murphy and D.W. Willis (eds.) Fisheries Techniques, 2nd edition. American Fisheries Society, Bethesda, Maryland.
- DeVries, D.A. and C.B. Grimes. 1997. Spatial and temporal variation in age and growth of king mackerel, *Scomberomorus cavalla*, 1977-1992. Fishery Bulletin. 95:694-708.
- Dutka-Gianelli, J., and D.J. Murie. 2001. Age and growth of sheepshead, *Archosargus probatocephalus* (Pisces: Sparidae), from the northwest coast of Florida. Bulletin of Marine Science. 68:69-83.
- Elsdon, T.S., Wells, B.K., Campana, S.E., Gillanders, B.M., Jones, C.M., Limburg, K.E., Secor, D.H., Thorrold, S.R., and Walther, B.D. 2008. Otolith chemistry to describe movements and life-history parameters of fishes - hypotheses, assumptions, limitations and inferences. Oceanography and Marine Biology: An Annual Review. 46:297-330.
- Escorriola, J.I. 1991. Age and growth of the gray triggerfish, *Balistes caprisкус*, from the southeastern United States. M.S. Thesis, University of North Carolina at Wilmington.

- Fable, W.A., A.G. Johnson, and L.E. Barger. 1987. Age and growth of Spanish mackerel, *Scomberomorus maculatus*, from Florida and the Gulf of Mexico. Fishery Bulletin. 85(4):777-783.
- Finucane, J.H., and L.A. Collins. 1986. Reproduction of Spanish mackerel, *Scomberomorus maculatus*, from the southeastern United States. Northeast Gulf Science 8(2):97-106.
- Finucane, J.H., L.A. Collins, H.A. Brusher, and C.H. Saloman. 1986. Reproductive biology of king mackerel, *Scomberomorus cavalla*, from the southeastern United States. Fishery Bulletin. 84(4):841-850.
- Fioramonti, C. Personal communication. NOAA Fisheries, Panama City Laboratory, Panama City, Florida.
- Fischer, A.J. Personal communication. Louisiana State University, Baton Rouge, Louisiana.
- Fischer, A.J. and B.A. Thompson. 2004. The age and growth of southern flounder, *Paralichthys lethostigma*, from Louisiana estuarine and offshore waters. Bulletin of Marine Science. 75(1):63-77.
- Fischer, A.J., M.S. Baker Jr., C.A. Wilson, and D.L. Nieland. 2005. Age, growth, mortality, and radiometric age validation of gray snapper (*Lutjanus griseus*) from Louisiana. Fishery Bulletin. 103:307-319.
- Fitzhugh, G. Personal communication. National Marine Fisheries Service, Panama City Laboratory, Panama City, Florida.
- Fore, P.L., and K.N. Baxter. 1972. Collection of larval Gulf menhaden, *Brevoortia patronus*, from Galveston Entrance (1959-1969) and Sabine Pass (1963-1967), Texas. U.S. Department of Commerce, Data Report 74 (COM-72-10584), 17 p.
- Friedland, K.D. and D.G. Reddin. 1994. Use of otolith morphology in stock discriminations of Atlantic salmon. Canadian Journal of Fisheries and Aquatic Sciences. 51:91-98.
- Gauldie, R.W., C.E. Thacker, West I.F., and L. Wang. 1998. Movement of water in fish otoliths. Comparative Biochemistry and Physiology (Part A). 120:551-556.
- González, P.D., S. Zupanovic and H.E. Ramis. 1974. Biología pesquera de la cherna americana del Banco de Campeche. INP/CIP Cuba. Res. Invest. 1:107-111.
- Grimes, C. B. 1978. Age, growth, and length-weight relationship of vermilion snapper, *Rhomboplites aurorubens* from North Carolina and South Carolina waters. Transactions of the American Fisheries Society. 107(3):454-456.

- Grimes, C.B., J.H. Finucane, L.A. Collins, and D.A. DeVries. 1990. Young king mackerel, *Scomberomorus cavalla*, in the Gulf of Mexico, a summary of the distribution and occurrence of larvae and juveniles, and spawning dates for Mexican juveniles. *Bulletin of Marine Science*. 46(3):640-654.
- Guindon, K. Personal communication. Florida Fish and Wildlife Conservation Commission, Florida Marine Research Institute, St. Petersburg, Florida.
- Heinke, F. 1905. Das Vorkommen und die Verbreitung der Eier, der Larven und der verschiedenen Altersstufen der Nutzfische in der Nordsee. Rapp. P.-V. Reun. Journal du Conseil Permanent International por l'Exploration de la Mer 3, Append. E, 41p.
- Hoffbauer, C. 1898. Die Altersbestimmung des Karpens an seiner Schuppe. *Allgemeine Fischereizeitung* 23(19).
- Holt, G.J., S.A. Holt, and C.R. Arnold. 1985. Diel periodicity of spawning in sciaenids. *Marine Ecology Progressive Series*. 27:1-7.
- Hood, P.B. and A.K. Johnson. 1997. A study of the age structure, growth, maturity schedules and fecundity of gray triggerfish (*Balistes capriscus*), red porgy (*Pagrus pagrus*), and Vermilion snapper (*Rhomboplites aurorubens*) from the eastern Gulf of Mexico. MARFIN Final Report.
- Hood, P.B., and R.A. Schlieder. 1992. Age, growth and reproduction of gag *Mycteroperca microlepis* (Pisces: Serranidae), in the eastern Gulf of Mexico. *Bulletin of Marine Science*. 51:337-352.
- Ibanez-Aguirre, A.L. and M. Gallardo-Cabello. 1996. Age determination of the grey mullet, *Mugil cephalus*, and the white mullet, *M. curema* in Tamiahua Lagoon, Veracruz. *Ciencias Marinas*. 22(3):329-345.
- Ihssen, P.E., H.E. Booke, J.M. Casselman, J.M. McGlade, N.R. Payne, and F.M. Utter. 1981. Stock identification: materials and methods. *Canadian Journal of Fisheries and Aquatic Sciences*. 38:1838-1855.
- Ingram, G.W., Jr. Personal communication. NOAA Fisheries, Pascagoula Laboratory, Pascagoula, Mississippi.
- Ingram, G.W., Jr. 2001. Stock structure of gray triggerfish, *Balistes capriscus*, on multiple spatial scales in the Gulf of Mexico. Ph.D. Dissertation, University of South Alabama. 228 p.
- Johnson, A.G. and C.H. Saloman. 1984. Age, growth, and mortality of gray triggerfish, *Balistes capriscus*, from the northeastern Gulf of Mexico. *Fishery Bulletin*. 82:485-492.
- Johnson, A.G. and L.A. Collins 1994 Age-size structure of red grouper, (*Epinephelus morio*), from the eastern Gulf of Mexico. *Northeast Gulf Science*. 13(2):101-106.

- Johnson, A.G., L.A. Collins and J. Jeffery Isely. 1993. Age-size structure of gag, *Mycteroperca microlepis*, from the northeastern Gulf of Mexico. *Northeast Gulf Science*. 13(1):59-63.
- Johnson, A.G., W.A. Fable, Jr., M.L. Williams, and L.E. Barger. 1983. Age, growth and mortality of king mackerel, *Scomberomorus cavalla*, from the southeastern United States. *Fishery Bulletin*. 81(1):97-106.
- Johnson, A.G., L.A. Collins, and J.J. Isely. 1998. Age-size structure of gag, *Mycteroperca microlepis*, from the northeastern Gulf of Mexico. *Northeast Gulf Science*. 13(1):59-63.
- Kalish, J.M, R.J. Beamish, E.B. Brothers, J.M. Casselman, R.I.C.C. Francis, H. Mosegaard, J. Panfili, E.D. Prince, R.E. Thresher, C.A. Wilson, and P.J. Wright. 1995. Glossary for Otolith Studies. Pages 723-729, *In: Recent Developments in Fish Otolith Research*. Belle W. Baruch Institute for Marine Biology and Coastal Research, Number 19, University of South Carolina Press, Columbia.
- Kimura, D.K. and J.J. Lyons 1991. Between-reader bias and variability in the age-determination process. *Fishery Bulletin*. 89:53-60.
- Kingsmill, S. 1993. Ear Stones Speak Volumes to Fish Researchers. *Science*. 260:1233-1234.
- Lee, R.M. 1920. A review of the methods of age and growth determination in fishes by means of scales. *Fishery Investigations - Series II (Sea Fisheries)*. 4(2):1-32.
- Limburg, K.E., R. Huang, and D. Bilderback. 2007. Fish otolith trace element maps: new approaches with synchrotron microbeam X-ray fluorescence. *X-ray Spectrometry*. 36:336-342.
- Lombardi-Carlson, L., G. Fithzugh, C. Palmer, C. Gardner, R. Farsky, and M. Ortiz. 2008. Regional size, age and growth differences in red grouper (*Epinephelus morio*) along the west coast of Florida. *Fisheries Research*. 91: 239-251.
- Lowerre-Barbieri, S.K., M.E. Chittenden, Jr., and C.M. Jones. 1994. A comparison of a validated otolith method to age weakfish, *Cynoscion regalis*, with the traditional scale method. *Fishery Bulletin*. 92:555-568.
- Lu, Z. and A.N. Popper. 1998. Morphological polarizations of sensory hair cells in the three otolith organs of a teleost fish: fluorescent labeling of ciliary bundles. *Hearing Research*. 126:47-57.
- MacNair, L.S., M.L. Domeier, and C.S.Y. Chun. 2001. Age, growth, and mortality of California halibut, *Paralichthys californicus*, along southern and central California. *Fishery Bulletin*. 99:588-600.

- Manooch III, C.S. and R. Matheson, III. 1981. Age, growth, and mortality of gray snapper collected from Florida waters. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies. 35:331-344.
- Mareska, J. Personal communication. Alabama Department of Conservation and Natural Resources, Alabama Marine Resources Division, Dauphin Island, Alabama.
- Markwitz, A., D. Grambole, F. Herrmann, W.J. Trompetter, T. Dioses, and R.W. Gauldie. 2000. Reliable micro-measurement of strontium is the key to cracking the life-history code in the fish otolith. Nuclear Instruments and Methods in Physics Research B-Beam Interactions with Materials and Atoms, 168:109-116.
- McErlean, A.J. 1963. A study of the age and growth of gag, *Mycteroperca microlepis* (Pisces: Serranidae), on the west coast of Florida. Florida Board of Conservation, Marine Laboratory Technical Series No. 41. 29 pp.
- Moe, M.A. Jr. 1969. Biology of the red grouper *Epinephelus morio* (Valenciennes) from the eastern Gulf of Mexico. Florida Department of Natural Resources, Marine Research Laboratory Professional Paper Series. 10:1-95.
- Moran, J. Personal communication. Atlantic States Marine Fisheries Commission, Washington D.C.
- Moran, P. and J. Baker. 2002. Polymerase chain reaction inhibition in historical fish scale mounting cards. Transactions of the American Fisheries Society. 131(1):109-119.
- Morison, A.K., S.G. Robertson, and D.G. Smith. 1998. An integrated system for production fish aging: image analysis and quality assurance. North American Journal of Fisheries Management. 18:587-598.
- Moyle, P.B. and J.J. Cech, Jr. 1988. Fishes: an introduction to ichthyology. Prentice-Hall, Inc. New Jersey. 559 p.
- Murie, D.J. Personal Communication. University of Florida, Fisheries and Aquatic Sciences, Gainesville, Florida.
- Murie, D.J., and D.C. Parkyn. 1999. Age, growth, and sexual maturity of white grunt in the eastern Gulf of Mexico: Part II. Final Report to S. Kennedy, Department of Environmental Protection, Florida Marine Research Institute, St. Petersburg, Florida 33701. 57 p.
- Murie, D. J., and D. C. Parkyn. 2008. Age, growth and sexual maturity of greater amberjack (*Seriola dumerili*) in the Gulf of Mexico. Final Technical Report for MARFIN NA05NMF4331071.
- Murphy, M.D. and R.G. Taylor. 1989. Reproduction and growth of black drum, *Pogonias cromis*, in Northeast Florida. Northeast Gulf Science. 10(2):127-137.

- Murphy, M.D. and R.G. Taylor. 1990. Reproduction, growth, and mortality of red drum *Sciaenops ocellatus* in Florida waters. Fishery Bulletin 88:531-542.
- Murphy, M.D. and R.G. Taylor. 1994. Age, growth, and mortality of spotted seatrout in Florida waters. Transactions of the American Fisheries Society. 123:482-497.
- Nicholson, W.R. 1978. Gulf menhaden, *Brevoortia patronus*, purse seine fishery: Catch, fishing activity, and age and size composition, 1964-73. U.S. Department of Commerce, NOAA Technical Report NMFS SSRF-722, 8 p.
- Nicholson, W.R., and W.E. Schaaf. 1978. Aging of gulf menhaden, *Brevoortia patronus*. Fishery Bulletin. 76(2):315-322.
- Ofori-Danson, P.K. 1989. Growth of grey triggerfish, *Balistes capriscus*, based on checks of the dorsal spine. Fishbyte (December):11-12.
- Old Dominion University/Virginia Marine Resources Commission. 2001. Glossary of Terms. Age and Growth Laboratory webpage (<http://web.odu.edu/sci/cqfe/age&growth/home.htm>).
- Palko, B.J. 1984. An evaluation of hard parts for age determination of pompano (*Trachinotus carolinus*), ladyfish (*Elops saurus*), crevalle jack (*Caranx hippos*), gulf flounder (*Paralichthys albigutta*), and southern flounder (*Paralichthys lethostigma*). NOAA Technical Memorandum NMFS-SEFC-132.
- Pannella, G. 1971. Fish otoliths. Daily growth layers and periodic patterns. Science. 173:1124-1127.
- Pannella, G. 1974. Otolith growth patterns: An aid in age determination in temperate and tropical fishes. Pages 28-39 In: T.B. Bagenal (ed) The Ageing of Fish. Unwin Brothers Ltd., Surrey, England.
- Pannella, G. 1980. Growth Patterns in Fish Sagittae. Pages 519-560, In: D.C. Rhoads and R.A. Lutz (eds) Skeletal Growth of Aquatic Organisms: Biological Records of Environmental Change, New York: Plenum Press.
- Pentilla, J, and L.M. Dery (eds). 1988. Age determination methods for Northwest Atlantic species. NOAA Technical Report NMFS 72.
- Petersen, C.G.J. 1892. Fiskeribiologiske forhold i Holboek Fjord, 1890-91. Beretning fra de Danske Biologiske Station for 1890(91), 1:121-183.
- Popper, A.N. and Z. Lu. 2000. Structure-function relationships in fish otolith organs. Fisheries Research. 46:15-25.

- Powell, D. 1975. Age, growth, and reproduction in Florida stocks of Spanish mackerel, *Scomberomorus maculatus*. Florida Department of Natural Resources, Florida Marine Research Publication No. 5, 21p.
- Prentice, J.A. and J.D. Wilfred, Jr. 1991. Use of known-age red drums to validate scale and otolith ages and to estimate growth in fresh water. *North American Journal of Fisheries Management*. 11:424-428.
- Radtke, R.L. 1989. Larval fish age, growth, and body shrinkage: information available from otoliths. *Canadian Journal of Fisheries and Aquatic Sciences*. 46:1884-1894.
- Radtke, R.L. 1990. Information storage capacity of otoliths: response to Neilson and Campana. *Canadian Journal of Fisheries and Aquatic Sciences*. 47:2463-2467.
- Radtke, R.L., D.W. Townsend, R.A. Kinzie, and D. Fey. 1999. Two-dimensional X-ray mapping of otoliths, a high-resolution technique for determining amphidromy in the tropical goby *Lentipes concolor* (Gill). *Journal of Experimental Marine Biology and Ecology*. 238, 21-27.
- Reibisch, J. 1899. Über die Eizahl bei *Pleuronectes platessa* und die Altersbestimmung dieser Form aus den Otolithen. *Wiss. Meeresuntersuchungen (Kiel) N.F.* 4:233-248.
- Richards, L.J., J.T. Schnute, A.R. Kronlund, and R.J. Beamish 1992. Statistical models for the analysis of aging error. *Canadian Journal of Fisheries and Aquatic Sciences*. 49:1801-1815.
- Richter, H. and J. G. McDermott. 1990. The staining of fish otoliths for age determination. *Journal of Fish Biology*. 36:773-779.
- Rodríguez, H. 1986 Estudio comparativo de dos estructuras rígidas (otolito y hueso mesopterigoide) para la estimación de edad crecimiento del mero (*Epinephelus morio*) del Banco de Campeche. Tesis Profesional. Univ. Autón., Nuevo León, México.
- Rohr, B.A. 1964. Comparison of the growth rings in the scales, otoliths, dorsal rays, and second anal spine as related to growth of the red drum, *Sciaenops ocellata*. *Journal of the Mississippi Academy of Sciences*. 10:208-212.
- Secor, D.H., J.M. Dean, and E.H. Laban. 1991. Manual for otolith removal and preparation for microstructural examination. Belle W. Baruch Institute for Marine Biology and Coastal Research. Technical Publication. 1991-01.
- Secor, D.H., J.M. Dean, and S.E. Campana. 1995a. Recent developments in fish otolith research. The Belle W. Baruch Library in Marine Science Number 19. University of South Carolina Press, Columbia.
- Secor, D.H., T.M. Trice, and H.T. Hornick. 1995b. Validation of otolith-based ageing and a comparison of otolith and scale-based ageing in mark-recaptured Chesapeake Bay striped bass, *Morone saxatilis*. *Fishery Bulletin*. 93:186-190.

- Sipe, A.M., and M.E. Chittenden, Jr. 2001. A comparison of calcified structures for aging summer flounder, *Paralichthys dentatus*. Fishery Bulletin. 99:628-640.
- Sinclair, M. 1988. Marine populations: an essay on population regulation and speciation. Washington Sea Grant Program. Seattle, WA.
- Sinclair, D.J., L.P.J. Kinsley, and M.T. McCulloch. 1998. High resolution analysis of trace elements in corals by laser ablation ICP-MS. Geochimica et Cosmochimica Acta. 62:1889-1901.
- Stanley, R.D. 1986. A comparison of age estimates derived from the surface and cross-section methods of otolith reading for Pacific ocean perch (*Sebastes alutus*). Proceedings of the International Rockfish Symposium (1986):187-196.
- Stevenson, D.K., and S.E. Campana. 1992. Otolith microstructure examination and analysis. Canadian Special Publication of Fisheries and Aquatic Sciences 117. 130 p.
- Stransky, C. 2001. Preliminary results of a shape analysis of redfish otoliths: comparison of areas and species. NAFO SCR Doc. 01/14, Ser. No. N4382. 9 p.
- Summerfelt, R.C. and G.E. Hall (eds). 1987. Age and Growth of Fish. Iowa State University Press. Ames, Iowa.
- Theiling, D.L. and H.A. Loyacano, Jr. 1976. Age and growth of red drum from a saltwater marsh impoundment in South Carolina. Transactions of the American Fisheries Society. 1:41-44.
- Thompson, B.A., J.H. Render, and R.L. Allen. 1989. Life History and Population Dynamics of Commercially Harvested Striped Mullet (*Mugil cephalus*) in Coastal Louisiana. Final Report FY 1987-1988 for Board of Regents', Rockefeller Fund Interest Earnings Grant Program. Louisiana State University, Baton Rouge, Louisiana. 83 p.
- Thompson, B.A., M. Beasley, and C.A. Wilson. 1999. Age distribution and growth of Greater Amberjack (*Seriola dumerili*) from the northcentral Gulf of Mexico. Fishery Bulletin. 97(2): 362-371.
- Tremain, D. Personal communication. Florida Fish and Wildlife Conservation Commission, Florida Marine Research Institute, St. Petersburg, Florida.
- Turner, W.R. 1969. Life history of menhadens in the eastern Gulf of Mexico. Transactions of the American Fisheries Society. 98(2):216-224.
- Valdés, E. and G. Padrón 1980 Pesquerías de palangre. Rev. Cub. Invest. Pesq. 5(2):38-52.
- Warren, J. Personal communication. University of Southern Mississippi, Institute of Marine Sciences, Gulf Coast Research Laboratory, Ocean Springs, Mississippi.

- Welsh, W.W. and C.M. Breder, Jr. 1924. Contributions to life histories of Sciaenidae of the eastern United States coast. *Bulletin of the U.S. Bureau of Fisheries*. 39:141-201.
- Wenner, C.A., W.A. Roumillat, J.E. Moran, Jr., M.B. Maddox, L.B. Daniel, III, and J.W. Smith. 1990. Investigations on the life history and population dynamics of marine recreational fishes in South Carolina: Part 1:2.1-2.35.
- Wilson, C.A. Personal communication. Louisiana State University, Baton Rouge, Louisiana.
- Wilson, C.A. and D.L. Nieland. 2001. Age and growth of red snapper, *Lutjanus campechanus*, from the northern Gulf of Mexico off Louisiana. *Fishery Bulletin*. 99:653-664.
- Wilson, C.A., J.B. Render, D.W. Beckman, and A.L. Stanley. 1988. The age structure and reproductive biology of sheepshead (*Archosargus probatocephalus*) landed in Louisiana. Final report of funded projects FY 1987-1988, Louisiana Board of Regents' Rockefeller Fund Interest Earnings Grant Program. Louisiana State University, Coastal Fisheries Institute, Baton Rouge, 49 p.
- Wilson, C.A., D.L. Nieland, and A.L. Stanley. 1995. Age, growth and reproductive biology of gray triggerfish (*Balistes capriscus*) from the northern Gulf of Mexico commercial harvest. Final Report, Coastal Fisheries Institute, Louisiana State University.
- Wischniowski, S.G., and S.J. Bobko. 2000. Age determination procedures for bluefish, *Pomatomus saltatrix*. Center for Quantitative Fisheries Ecology, Old Dominion University, Norfolk, Virginia. 19 p.
- Zhao, B., J. C. McGovern, and P. J. Harris. 1997. Age, growth, and temporal change in size-at-age of the vermilion snapper from the South Atlantic Bight, based on otolith sections. *Transactions of the American Fisheries Society*. 126:181-193.

7.0 Glossary of Terms Used in Age and Growth Studies

Sources used to compile this glossary include: Summerfelt and Hall 1987, Secor et al. 1991, Kalish et al. 1995, C.A.R.E. 1997, Old Dominion University /Virginia Marine Resources Commission 2001.

A

accuracy - the closeness of a measure or computed value to its true value.

age - a unit to express the passage of time to capture measured in years, months, days or other units.

age-group (age-class, cohort-age) - a group of fish that have the same assigned age within a given time period (e.g., five-year-old age-group); the term is not synonymous with year-class.

age estimation, age determination - the preferred terms for the process of assigning ages to fish as opposed to the term aging (ageing), which refers to time-related processes such as the alteration of an organism's composition, structure, and function.

ampulla - the enlarged chamber containing a patch of sensory epithelium at one end of each semicircular canal of the inner ear.

annual age - an integer enumeration of age corresponding to year-class.

annual growth zone - all growth on a structure which forms during one year; consisting of an opaque zone or annulus and a translucent zone, generally formed during the winter and summer months, respectively.

annulus (pl. annuli) - a continuous, concentric growth zone that forms once a year, for most fish during a period of slow or no growth (see opaque growth zone, winter growth zone); the optical appearance of these marks depends on the otolith structure and the species.

anterostrum - an anterior projection of the sagitta located dorsal to the sulcus acousticus and rostrum; generally shorter than the rostrum.

aragonite - an inorganic, crystalline polymorph of calcium carbonate that combines with otolin to form the otolith matrix.

asteriscus (pl. asterisci) - one of the three otolith pairs

found in the membranous labyrinth of osteichthyan fishes; lies within the lagena of the pars inferior.

B

biological age - the time elapsed from estimated birth to capture expressed in years and fractions of years.

birth date (theoretical) - calendar date that coincides with the mode of spawning activity for a given species.

blind reading - visual assessment of otolith annuli and margin/edge development with no knowledge of fish size and date of capture.

C

calendar age - the age of a fish based on a calendar year rather than to the true date of hatching.

calendar birthdate - January 1; used to maintain year classes when biological birthdate is unknown.

cauda - the posterior, medial-extending section of the sulcus acousticus.

check - a discontinuity (e.g., a stress-induced mark) that forms within the translucent zone, denoting a slowing of growth; checks do not form annually but reflect various environmental or physiological changes; distinguished by the width of the zone relative to annuli, location relative to annuli, and incomplete formation or poor definition.

circuli (circulus; singular) - fine ridges laid in a circular pattern around the focus of a scale.

cohort - group of fish that begins life about the same time and is produced during a relatively discrete spawning event; difficult to apply to fishes that spawn monthly or some other periodicity; does not imply year-class.

cohort age - see annual age.

core - the primordium of the otolith (sometimes used synonymously with focus).

core region - the area or areas surrounding one or more primordium.

corroboration - a measure of the consistency or repeatability of an age determination method when two different readers agree on the number of zones present; not to be confused with validation.

crystallized otolith - an otolith displaying inadequate calcification; age determinations are generally not possible due to missing annuli.

D

daily increment - an increment formed over a 24-hour period; synonymous with daily growth increment and daily ring.

distal edge - the external margin of an otolith cross-section.

distal surface - the external surface of a whole otolith; the surface opposite the sulcus.

E

edge type - synonymous with edge zone; extent of opaque or translucent deposition on the outer margin of the otolith representing the most recent growth.

F

false annulus (pseudoannulus) - sometimes used synonymously with “check” or “check mark;” refers to a zone of slow growth that is not a true annulus; also, a characteristic ring on otoliths that occurs before the first annulus and fairly close to the nucleus.

focus - the hypothetical or real point of origin of an otolith or scale; the starting point of a sectioned or whole otolith where the reader chooses to start a count or use as a reference point for measurement.

I

increment - the region between similar zones on a structure used for age estimation; the term refers to a structure, but it may be qualified to refer to portions of the otolith formed over a specified time interval (e.g., subdaily, daily, annual); an annual increment is made up of an opaque zone and a translucent zone, whereas a daily increment consists of a D-zone and an L-zone.

L

lagena - an organ of non-mammalian vertebrates analogous to the cochlea.

lapillus (pl. lapilli) - one of the three otolith pairs found in the membranous labyrinth of osteichthyan fishes; lies within the utriculus of the pars superior.

M

marginal increment - the region beyond the last identifiable estimation mark at the otolith margin; usually expressed in relative rather than quantitative terms, i.e., as a fraction or proportion of the last complete increment; see edge type.

N

nucleus - central portion of an otolith; used synonymously with core, focus, kernel, or primordium.

O

opaque growth zone - usually synonymous with winter growth zone; a banded region of an otolith section that interferes with the passage of transmitted light and therefore appears dark relative to adjacent translucent growth zone(s); appears bright under reflected light; usually an area of high concentrations of calcium aragonite relative to otolin; occasionally, opaque zones are formed in areas where the aragonite crystal alignment interferes with light transmission through the otolith section; the opaque and translucent growth zones together form the annual growth zone.

ostium - the anterior section of the sulcus acusticus.

otolin - the organic protein found in the otolith, closely related to conchiolin of some mollusks.

P

precision - the closeness of repeated measurements of the same quantity; in age determination, it relates to the variability between or within readers.

primordium (pl. primordia) - the initial deposition site of organic matrix and calcium carbonate of an otolith; if several primordia are present, they generally fuse to form the otolith core.

proximal edge - the internal margin of an otolith cross-section.

proximal surface - the internal surface of a whole otolith; the surface on which the sulcus is found.

R

radii (radius; singular) - linear extensions of ridges from the focus to the anterior margin of a scale.

reading axis - preferred path along which annuli are counted; see sulcus edge.

ring (band, zone, check) - a descriptive term used in determining the age of a fish from hard parts; does not necessarily designate yearly or annual marks.

rostrum - anterior-most, ventral projection of the sagitta; generally longer than the anterostrum.

S

sacculus - the smaller chamber of the membranous labyrinth of the inner ear.

sagittae (sing. sagitta) - largest of three otolith pairs within the membranous labyrinth of osteichthyan fishes and therefore most often selected for otolith studies; lies within the sacculus of the pars inferior; generally compressed laterally and elliptical in shape with wide variation in appearance among species.

semicircular canal - any of the loop-shaped tubular parts of the labyrinth of the inner ear that together constitute a sensory organ consisting of an inner membranous canal and a corresponding outer bony canal formed in a group of three in planes nearly at right angles to each other.

split - discontinuity in an annular zone, analogous to a check; causes the annulus to appear as two or more closely spaced winter zones.

subdaily increments - an increment formed over a period of less than 24 hours.

sulcus acusticus/acousticus - commonly called sulcus or sulcus groove; a longitudinal sculptured groove extending down the convex (medial) surface of a sagittal otolith through which an auditory nerve passes; frequently referred to in otolith work because of the clarity of increments near the sulcus in transverse sections of sagittae.

sulcus edge - on an otolith cross-section, the margin adjacent to the sulcus on the internal or proximal surface.

summer growth zone - see translucent growth zone.

T

transition zone - a marked change in the annual growth zone requiring an adjustment to age-reading criteria.

translucent growth zone - the banded regions on an otolith section that allow a greater passage of transmitted light relative to the opaque or winter zones; usually an area of high concentrations of otolin relative to calcium aragonite; represents a period of faster growth; also called summer zone; the term hyaline has been used, but translucent is the preferred term.

U

utricle - the part of the membranous labyrinth of the inner ear into which the semicircular canals open.

V

validation - the process of proving that otolith rings accurately represent annual growth patterns which can be used to assign an age to a fish; methodologies include tag and recapture, hatchery releases, and chemical or temperature marking of otoliths.

verification - the process of determining ageing precision comparing ages assigned blindly by multiple readers.

W

winter growth zone - see opaque growth zone; represents a period of slower growth.

Y

year class - fish spawned or hatched in a given year.

8.0 Appendices

8.1 Permanent equipment and apparatus	8-2
8.2 Expendables	8-4
8.3 Contact information for Appendix 8.1 and 8.2	8-9
8.4 Photo credits	8-11

Appendix 8.1 Suppliers and Supplies for Otolith Processing - Permanent Equipment and Apparatus

Item Description	Purpose	Manufacturer or Supplier	Model or Stock No.	Count per Unit	Current Price
Small hotplate, 120v	Warming thermoplastic adhesive or araldite resin	Barnstead/Thermolyne	HP131225Q	1	213.00
Motorized polishing apparatus	Polishing otolith sections	Fisher Scientific	C5505070 Crystal Master Plus 6 and Master Lap 115V	1	204.00
Buehler polishing machine	Polishing otolith sections	Buehler	EcoMet® 250 without powerhead 49-7200	1	4,995.00
Top-loading balance	Gravimetric measurement of embedding epoxy resin and hardener components	Lab Safety Supply	12002 ACCULAB 120 gram	1	387.00
Low-speed diamond wafering saw	Cutting otolith thin sections	Buehler, Ltd.	Isomet Model 11-1280-160 Low Speed Saw	1	4,595.00
Low-speed diamond wafering saw	Cutting otolith thin sections	Struers	Minitom	1	3,750.00
High-speed diamond wafering saw	Cutting otolith thin sections	Struers	Accutom-5	1	18,000.00
Low-speed diamond wafering saw, 0-300rpm	Cutting otolith thin sections	South Bay Technology	Model 650	1	Call for price
Low-speed diamond wafering saw, 0-300rpm	Cutting otolith thin sections	Buehler, Ltd.	Model 11-1280-160	1	Call for price
High-speed diamond wafering saw, 0-1,000rpm	Cutting otolith thin sections	Buehler, Ltd.	Model 11-2810	1	Call for price
Thin section machine	Cutting otolith thin sections	Hillquist	Model 1005	1	6,400.00
Stereo-microscope with transmitted and reflected light sources	Use with or without imaging system to read otolith annuli and edge zones	Wild-Leitz, Olympus, Parco, Leica, Meiji, etc.	---	1	Call for price

Item Description	Purpose	Manufacturer or Supplier	Model or Stock No.	Count per Unit	Current Price
Compound microscope	Use with or without imaging system to read otolith annuli and edge zones	Wild-Leitz, Olympus, Parco, Meiji, etc.	---	1	Call for price
Optimas color optical image analysis software	Otolith image capture, enhancement, and display; spatial measurement of growth zones	Meyer Instruments Optimas Corp.	Contact for regional sales representative and technical support		Call for price
Image-Pro Plus image analysis software	Otolith image capture, enhancement, and display; spatial measurement of growth zones	Media Cybernetics	Contact for regional sales representative and technical support		Call for price
Low-power dissecting microscope	Judging quality of otolith sections	Wild-Leitz, Olympus, Parco, Meiji, etc.	---	1	Call vendor for price quote
Incubator oven	Curing resin for embedded otoliths	Fisher Scientific	.7 cu. ft. 13-246-506		928.00
UV lamp unit	Used to cure Loc-Tite adhesive on otolith section slide mounts	Ultra-Violet Products (UVP)	EL Series, Model UVL-28	1	Call for price
Cooling fan for UV lamp cabinet enclosure		RadioShack	120v AC 4-in. fan	1	24.99
Slide storage cabinet	Used to stack microscope slide trays during sample processing	Fisher Scientific	Catalog No. 12-580	1	646.68
Aluminum slide tray, 20-slide capacity	Used to hold slides during sample processing	Fisher Scientific Catalog	No. 50-931-20	1	140.64

Appendix 8.2 Suppliers and Supplies for Otolith Processing - Consumables

Item Description	Purpose	Manufacturer or Supplier	Model or Stock No.	Count per Unit	Current Price
Sample envelopes	Field collection and permanent archiving of whole otoliths	Allometrics	VW 56775-039 3 x 5 brown Kraft tin-tie safety fold	250/box	24.50
Plastic sample vials (20ml)	Field collection and permanent archiving of whole otoliths	Wildlife Supply Co.	Gummed, 2.25X3.5 inch 114-B20	1000/box	125.00
Scintillation vials with caps (20ml)	Field collection and permanent archiving of whole otoliths	Fisher Scientific	03-341-71A	100/case	316.03
Micro-centrifuge tubes with caps	Field collection and permanent archiving of whole otoliths	Lab Safety Supply	107279 Glass	100/box	38.40
Tissue culture trays, 24 well	Field collection of small otoliths	VWR Scientific	20170-610	500/bag 10 bags/case	189.00
Tissue culture trays, 12 well	Storing/Archiving of whole otoliths and otolith sections	VWR Scientific	29443-952	50/case	157.02
Polyethylene embedding molds, 22 x 30mm	Storing/Archiving of whole otoliths and otolith sections	VWR Scientific	29442-040	50/case	143.79
Flat embedding mold, reusable	Embedding otoliths for thin sectioning	VWR Scientific Polysciences	15160-270 Peel-Away embedding molds	8 cells/tray 36 trays/case	53.00
Histological disposable base molds	Embedding very small otoliths for thin sectioning and whole larval fish for grinding	Ted Pella Polysciences	105 Pelco 24-cavity, 3mm x 14mm x 7mm 10504 (clear)	1 1	11.45 10.75
	Embedding otoliths for thin sectioning	Surgipath Medical Industries		500/case	50.00-60.00

Item Description	Purpose	Manufacturer or Supplier	Model or Stock No.	Count per Unit	Current Price
Clear silicon embedding molds	Embedding otoliths for thin sectioning	Electron Microscopy Sciences	70900	1	8.50
Araldite resin and hardener	Embedding otoliths for thin sectioning	Electron Microscopy Sciences	Araldite GY 502 (resin) and Aradur 956-1 US (hardener)	450ml	14.00
Epoxy resin and hardener	Embedding otoliths for thin sectioning	Ribelin Sales, Inc	Epoxy resin 7132	50-lb. pail	Call for price
Disposable plastic beakers, 100 ml	Mixing two-part embedding epoxy	Artstuf	Epoxy hardener 2001	8 lbs	47.85
Loctite Hysol 0151 epoxy	Embedding otoliths for thin sectioning	VWR Scientific	13915-624 Tri-Pour	4 lbs	43.70
Latex gloves	Skin protection while mixing and pouring embedding media	Motion Industries	Hysol 151 parts A and B in kit	100/box	29.00
Nitrile gloves	Skin protection while mixing and pouring embedding media	Ellsworth Adhesives	15 W 1071	2.6 lb kit 3/pack	200.12
Magni-Visor	Close-up work: cleaning otoliths, positioning in embedding molds, marking epoxy blocks for sectioning	Ward's	19-130-1597	100/box	18.00
Metal spatulas	Mixing embedding epoxy	Fisher Scientific	25 W 2101	1	25.00
Thin Section Machine Diamond Blade	Cut-off Wheel	Hillquist	15 W 4313	12/pack	23.00
Thin Section Machine Diamond Cup Wheel	Precision Grinding	Hillquist	8 inch	1	180.00
Thin Section Machine Diamond Blade	Cut-off Wheel	Hillquist	8 inch	1	665.00
		Hillquist	HCR-100 Diamond Blade, 8 x .050 x 5/8	1	165.00

Item Description	Purpose	Manufacturer or Supplier	Model or Stock No.	Count per Unit	Current Price
Diamond wafering blade	High-speed saw	Precision Surfaces International	4 inch 6 inch	1 1	177.00 300.00
Diamond wafering blade	Low-speed saw	Buehler, Ltd.	11-4244 Isomet High Concentration Wafering Blade	1	227.00
Diamond wafering blade	Low-speed saw	Struers	230CA Diamond cut-off wheel	1	200.00
Diamond wafering blade	Low speed saw	South Bay Technology	DWH4122 Diamond wheel - 4" x .012" x medium/high concentration	1	call for price
Diamond wafering blade	Low-speed saw	Alro Industrial Supply	Norton Diamond Wheel, Product No. JO-588-678	1	99.00
Diamond wafering blade	Low-speed saw	Diamond Wheel, Inc	ME120928	1	96.00
Diamond wafering blade	Low-speed and high-speed diamond wafering saws	Struers Inc.	Diamond Cut-off Wheel MOD10	1	289.00
Glass microscope slides, 1.2 mm thickness	Otolith mounts	Ward's	14 W 3501	72/pack 10 packs/box	85.00
Thermoplastic mounting adhesive	Securing otolith thin sections and whole otoliths on glass slide mounts	Aremco Products	Crystalbond 509	5 sticks/pack	70.00
Thermoplastic mounting adhesive	Securing otolith thin sections and whole otoliths on glass slide mounts	Hugh Courtright & Co.	Lakeside Brand (Quartz) Thermoplastic Cement, Stock No. 70C	12 bars/box	33.45
Loctite mounting adhesive	Securing otolith thin sections and whole otoliths on glass slide mounts	Loctite Corp. Motion Industries	Loctite 349Impruv® Light Cure Adhesive (UV activated)	1 1	23.67

Item Description	Purpose	Manufacturer or Supplier	Model or Stock No.	Count per Unit	Current Price
Cytoseal Mounting Media	Covering and securing thin sections on glass mount slides	Cole Parmer	EW-48802-00 (16 oz)	1	272.00
Flo-Texx	Covering and securing thin sections on glass mount slides	Fisher Scientific	143904 143903	4 pack 6 pack	272.99 177.93
Micropolishing compound, .3 micron	Polishing apparatus	Buehler, Ltd.	40-6363-006 Alumina II	6 oz.	16.00
Wet-dry sandpaper, 600 grit	Polishing apparatus	Hardware store	---	multi-sheet pack	10.00
Polishing cloth	Polishing otolith thin sections	Buehler, Ltd.	MicroCloth®, PSA backed 40-7216 (6 inch disc) 40-7218 (8 inch disc)	10/pack 10/pack	33.00 40.00
Disposable glass pipettes, 10 ml	Dispensing epoxy into small molds (e.g., those used for larval fish otoliths) during otolith embedding	Ward's	17 W 1308	1	7.00
Blunt-fine tipped scissors	Trimming otolith sections prior to mounting	Ward's	14 W 0940	1	5.00
Kimwipes	Wiping saw coolant from otolith sections	Lab Safety Supply	14011 Extra Low Lint, Delicate Task, 11" X 17"	140/box	6.00
Teri-Towels	General housekeeping	Ward's	15 W 1024		17.00
Water soluble saw coolant	Diamond saw blade lubricant and coolant	South Bay Technology	02-02460	1 qt. concentrate	---
Soluble oil	Diamond saw blade lubricant and coolant	Buehler, Ltd.		1 gallon	50.00
Glycerine		Fisher Scientific	G33-4	4 bottles/case	305.33

Item Description	Purpose	Manufacturer or Supplier	Model or Stock No.	Count per Unit	Current Price
Plastic slide boxes	Storing slide	Fisher Scientific	03448-5	72 boxes/case	565.25
Plastic slide boxes	Storing slide	Fisher Scientific	82003-412	72 boxes/case	1,080.97
Plain glass slides	Mounting otolith sections	Fisher Scientific	12-550A	10 gross/case	136.81
Frosted glass slides	Mounting otolith sections	Fisher Scientific	12-550-43	10 gross/case	175.66
Disposable droppers	Dispensing embedding and mounting media	Fisher Scientific	6219-0068	100 pack	50.03
Rubber cement	Securing polishing cloth and wet-dry sandpaper to polishing apparatus	Office supply	4 oz	1	1.59
Forceps	Handling otolith thin sections and otolith mounts	Fisher Scientific	08-953E straight 08-953I curved	1 1	46.42 67.45
Ultraviolet fluorescent light bulb (24")	curing of Loctite adhesive	General Electric	F20T12BLB	1	40.00
Stir bar	Mixing Araldite resin and hardener	Fisher Scientific	14-512-143	1	2.98
Plastic slide boxes	Long term slide storage	Ted Pella	2102-1 100 slide capacity	72 boxes/case	388.80
Hot melt glue gun	Mounting whole otoliths to mounting paper for production sectioning	local hardware/craft store		1	15.00
Hot melt glue sticks	Mounting whole otoliths to mounting paper for production sectioning	local hardware/craft store		24 pack	5.30
Food service trays, various sizes	Platform for embedding molds during pouring of epoxy	Science supply or restaurant supply			

Appendix 8.3 Contact information for Appendix 8.1 and 8.2

Allometrics, Inc.
PO Box 15825
Baton Rouge, LA 70895
(800) 528-2246
www.allometrics.com

Alro Industrial Supply
12490 49th Street
Clearwater, FL 34622-4310
www.alro.com

Aremco Products, Inc.
PO Box 429
Ossining, NY 10562
(914) 762-0685
www.aremco.com

Barnstead/Thermolyne
PO Box 797
2555 Kerper Boulevard
Dubuque, IA 52004-0797
(319) 556-2241
www.barnsteadthermolyne.com

Buehler, Ltd.
41 Waukegan Road
Lake Bluff, IL 60044
(800) 283-4537
www.buehlerltd.com

C & H Distributors
770 S. 70th Street
PO Box 14770
Milwaukee, WI 53214
(414) 443-1700
www.chdist.com

Ciba-Geigy Corporation
Formulated Systems Group
4917 Dawn Avenue
East Lansing, MI 48823
(800) 875-1363

Crystalite Corporation
8499 Green Meadows Drive
Westerville, OH 43081
(800) 777-2894

Diamond Wheel, Inc.
440 Union Place
Excelsior, MN 55331
(800) 328-0303
diamondwheelinc.com/

Electron Microscopy Sciences
PO Box 251
321 Morris Road
Fort Washington, PA 19034
(800) 523-5874

Fine Science Tools
1500 Industrial Way
Belmont, CA 94002
(800) 521-2109

Fisher Scientific
2775 Pacific Drive
PO Box 4829
Norcross, GA

Hillquist, Inc.
35502 S.E. Fall City Snoqualmie Road
Fall City, WA 98024
(425) 222-6968
www.hillquist.com

Hugh Courtright & Co., Ltd.
4314 West 166th Street
Oak Forest, IL 60452
www.right-tape.com

Lab Safety Supply
PO Box 610 Vineland, NJ 08360
(800) 356-0783
www.labsafety.com

Loctite Corporation
1001 Trout Brook Crossing
Rocky Hill, CT 06067
www.loctite.com

Martin Microscope Company
207 South Pendleton Street
Easley, SC 29640
www.martinmicroscope.com

Meyer Instruments
1304 Langham Creek, Suite 235
Houston, TX 77084
(281) 579-0342
www.meyerinst.com

Motion Industries
(225) 356-6131
www.motion-industries.com

National Diagnostics
305 Patton Drive
Atlanta, GA 30336
(800) 536-3867

Optimas Corporation
19811 North Creek Parkway
Bothell, WA 98011
(800) 635-7226
www.optimas.com

Polysciences, Inc.
Corporate Headquarters
400 Valley Road
Warrington, PA 18976
(800) 523-2575
www.polysciences.com

Precision Surfaces International
922 Ashland Street
Houston, TX 77008-6734
(713) 426-2220
(800) 843-0950

South Bay Technology, Inc.
1120 Via Callejon
San Clemente, CA 92672
(714) 492-1499
www.southbaytech.com

Struers, Inc.
810 Sharon Drive
Westlake, OH 44145
1-888-787-8377
www.struers.com

Surgipath Medical Industries, Inc.
PO Box 528
Richmond IL 60071
(800) 225-3035
www.surgipath.com

Ted Pella, Inc.
PO Box 492477
Redding, CA
(800) 237-3526
www.tedpella.com

VWR Scientific Products
(800) 932-5000
www.vwrsp.com

Ward's Natural Science Establishment, Inc.
(800) 962-2660
www.wardsci.com

Appendix 8.4 Photo and Illustration Credits

Andy Fischer	3.22, 3.23, 3.24, 3.25
Joseph Smith	5.125, 5.126, 5.127, 5.128, 5.129
Britt Bumgardner	2.4, 2.5, 3.6, 3.9, 3.10, 3.11, 5.11, 5.14, 5.22
Debra Murie	3.28, 3.29, 3.42, 3.43, 3.44, 3.45, 3.52, 5.74
Chris Palmer	3.27, 4.10, 5.65, 5.80
John Mareska	3.37, 3.39, 5.27, 5.28, 5.29, 5.30, 5.31, 5.32
Stacey Randall	5.2, 5.10, 5.18, 5.26, 5.35, 5.42, 5.51, 5.60, 5.68, 5.77, 5.83, 5.91, 5.98, 5.105, 5.112, 5.119
Luiz Barbieri	5.87
Walter Ingram	3.36, 5.43, 5.46, 5.47, 5.48
Kristin Maki (VIMS)	3.34
Rich McBride	3.30
Janet Tunnell	3.31, 3.32, 3.33
Jim Franks	3.35
Carrie Fioramonti	5.41, 5.44, 5.45
Jessica Carroll	3.19, 3.20, 3.21
Tut Warren	4.2, 4.3, 4.6, 4.8, 4.12, 4.14
Ivy Baremore	3.46, 3.47, 3.48
Daniel Merryman	3.50
Stephan Wischnowski	3.57
NOAA Fisheries/ Panama City Laboratory	5.55, 5.57, 5.67, 5.69, 5.70, 5.102, 5.115, 5.122, 5.124
FWC/FWRI	All Diane Peebles fish illustrations in Section 5 provided with permission by FWC (Southern Flounder image modified by S. VanderKooy)

All other images and illustrations by S. VanderKooy, J. Rester, and R. Hode unless otherwise noted

**Gulf States Marine Fisheries Commission
2404 Government Street
Ocean Springs, Mississippi, 39564**

